

1

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

REMUS

Fungicide

Wettable Powder

75% Chlorothalonil

(2,4,5,6-tetrachloro-1,3-benzenedicarbonitrile)

Acute oral in rats LD₅₀ > 5000 mg/kg

CAUTION

READ THE LABEL FIRST

DO NOT USE AT HOME

KEEP AWAY FROM CHILDREN AND FOOD

DO NOT INHALE THE PRODUCT VAPOR AND ITS DRIBLETS

WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES

DO NOT EAT DURING THE APPLICATION, DO NOT SMOKE

AVOID SKIN AND EYE CONTACT

HUMANS AND ANIMALS FOR 3 DAYS ARE NOT ALLOWED INTO THE AREA APPLIED

THE USE ON THE PRODUCTS OTHER THAN RECOMMENDED STRICTLY PROHIBITED

PROTECTION PREVENTIONS: Although Remus has low acute, subacute and oral, dermal toxicity, the regular protection preventions must be taken in case applying it. While spraying, the driblets should not be inhaled, eye and skin contact must be avoided, and after use, hands and face must be washed well with water and soap. Never leave the patient alone. In case of eye contact, immediately wash well with water during min. 15 minutes. The product has temporary allergic side effects such as eruption on open skin areas or pink eyes in case of contact, slight irritation on bronches. Has no permanent effects. The patients with allergic metabolisms must see a doctor in such cases, must be taken to doctor if swallowed with the package and the label as well.

ANTIDOTE :No specific antidotes. Treatment according to symptoms.

NPCA (NATIONAL TOXIC CONSULTANCY ASSOCIATION) Tel:114

2

Production date:

Expiry date:

Charge nr:

License date and nr:30.12.2009

THE PLANTS AND DISEASES TO APPLY ON

| Plant | Disease | Application Dose and Term | Period Between Last Spraying and Harvest(days) |
|-----------|----------------------------|---------------------------|--|
| Apple | Venturia inaequales | 150 g/100 l water | 3 days |
| Pear | Venturia pirina | 200 g/100 l water | 14 days |
| Tomato | Phytophthora infestans | 200 g/100 l water | 3 days |
| Peanuts | Aspergillus niger | 500 g/100 kg seed | ----- |
| Chickpeas | Mycosphaerella rabiei | 200 g/100 l water | 3 days |
| Sugarbeet | Cercospora beticola | 200 g /da | ----- |
| Cucumber | Pseudoperonospora cubensis | 200 g/da | 3 days |

PREPARING TO APPLY:

Fill the half of the tank with water. The necessary amount of the product is mixed with a little amount of water in a separate tank and added into. The water left is added while the mixer of the apparel is on. In case of spraying the plants must be washed in the best way, under foil nozzles must be applied onto the ground tools, so that the back surfaces of the leaves can also be wetted. The spraying must be on the days that are not windy.

Venturia inaequales:

- 1.spraying, by the time the flower spots pop.
2. spraying at the pink badge bud term.
- 3.spraying by the time 70-80% of the petals fall
4. and the rest, according to the weather conditions and the development of the disease.

Venturia pirina:

- 1.spraying, by the time the flower spots pop.
2. spraying at the white badge bud term.
- 3.spraying by the time 70-80% of the petals fall
4. and the rest, according to the weather conditions and the development of the disease.

Phytophthora infestans:

Spraying starts by the time the prior marks of 3-5 mm width are noticed on the tomato leaves.

Mycosphaerella rabiei:

Spraying starts by the time round marks of 2-3 mm width or capsule shaped marks, and are repeated with intervals of 7-10 days according to the development of the disease and the meteorologic conditions.

Cercospora beticola:

The sprayings are started by the time the prior symptoms are noticed and repeated every 20 days.

MISCIBILITY:

Before Remus is mixed with the wettable in water powder formulations of many kinds of insecticides, fungicides, and acaricides, its miscibility and its state of phytotoxicity must be tested. Must never be mixed with products that have very strong acid and alkali characteristic.

3**Net Amount:****Gross Amount:****Max. Retail Sales Price:**

*Toxic for bees.

*Toxic for the fish, do not contaminate into the waters.

*Keep the product in its package closed.

*Exterminate the emptied bottles accordingly and do not use for any other purpose.

*Keep away from its powder while applying the product.

CAUTIONS ON USE AND STORAGE:

If stored under normal storing conditions, at cool and dry places, in the original package, the product keeps its physical, chemical, and biological properties max. for two years within tolerance limits.

COMPANY'S ANNOUNCEMENT:

The company guarantees the quality of the product provided that it is stored in normal conditions and in the original package and sold in the original and closed package, but denies the responsibility for the damages that may result of applying the product in a wrong way or irregularly by not following the advices, or storing and applying in a wrong way.

MANUFACTURER and LICENSED COMPANY:

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79