

Manufacture Date :
Expiration Date :
Charge No :
Permit Date & Number :

TARGETED PLANTS AND PESTS

PLANT NAME	PEST NAME	DOSAGE & TIME PERIOD
WHEAT	Bifora (<i>Bifora radians</i>)	100 ml/da at wheat tillering period, in 2-4 leaved period of bifora.
SUGAR BEETS	Bifora (<i>Bifora radians</i>)	100 ml/da after beet and weeds emerge, and in 2-4 leaved period of weeds.
	Creeping Thistle (<i>Cirsium arvense</i>)	
	Milkweed (<i>Sonchus oleraceus</i>)	
	Eastern groundsel (<i>Senecio vernalis</i>)	
	Camomile (<i>Matricaria</i> spp.)	
	Nodding thistle (<i>Carduus nutans</i>)	
	Prickly Lettuce (<i>Lactuca scariola</i>)	
Cornflower (<i>Centaurea cyanus</i>)		

METHOD OF APPLICATION:

Preparation:

Korazim readily mixes with water and therefore it can be used with any type of applicator. Fill the applicator tank with water to the half point. Mix the necessary amount of product with water in a separate container and pur this preparation into the tank. Keep tank mixer running and add water until the tank is full.

Calibration of Applicator:

Applicator should be calibrated to ensure correct and homogenous dispersal of the recommended dosage per decare. To do this first check that all nozzles of the applicator are functional and sprays same amount of water per unit of time by placing containers under each nozzle and making sure the amount of water gathered in each container is equal. Fill the applicator tank with plain water. Spray a certain area with a consistent speed as in pesticide application. Then measure the sprayed area and amount of consumed water to calculate the amount of product necessary for one tank full of water. Prepare the product as explained above and pour into the tank. Then proceed to spraying with the same speed and rate as the calibration application. It is applied with 20-40 liter water per decare on recommended dosages.

Signature and Stamp
Dr. Yunus BAYRAM
Department Chief

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



KORAZIM

Soluble Concentrate (SL)

Name & Amount of Active Ingredient: 100 g
Clopyralid in Liter
Clopyralid (3.6-Dichloropicolinic acid) 3.6-
dichloropyridine-2-carboxylic acid

GROUP	O	HERBICIDE
-------	---	-----------

FIRSTLY READ THE LABEL.
DO NOT USE AT HOME.
KEEP AWAY FROM CHILDREN, FOODSTUFFS AND ANIMAL FEED.
DO NOT INHALE VAPORS AND/OR PARTICLES OF PESTICIDE.
PLEASE USE PROTECTIVE CLOTHING, GLOVES AND GOGGLES.
DO NOT EAT, DRINK OR SMOKE DURING APPLICATION OF PESTICIDE.
DO NOT ALLOW HUMAN AND ANIMAL TO ENTRY INTO PESTICIDE TREATED AREAS FOR 1 DAY
PLEASE FOLLOW THE USAGE INSTRUCTIONS TO PREVENT ANY HUMAN HEALTH AND ENVIRONMENTAL RISKS.
IT IS STRICTLY PROHIBITED TO USE THE PESTICIDE ON ANYTHING OTHER THAN RECOMMENDED TARGETS.
SHAKE WELL AFTER ADDING CLEAN WATER TO EMPTY CONTAINER OF PESTICIDE WITH THE RATE OF 1/4. POUR THE SHAKEN WATER INTO DISINFECTION TANK. REPEAT THIS PROCESS THREE TIMES

**NATIONAL POISON INFORMATION CENTRE (UZEM)
PHONE: 114**

5 December 2014
MINISTRY OF FOOD AGRICULTURE AND LIVESTOCK
GENERAL DIRECTORATE OF FOOD AND CONTROL
ACCEPTED

Net Amount :
Max. Retail Sales Price :

HAZARD TYPE:

Xi: IRRITANT
N: DANGEROUS FOR THE ENVIRONMENT

HAZARD SYMBOLS:



Xi: IRRITANT **N:** DANGEROUS FOR THE ENVIRONMENT

INGREDIENTS:

Contains clopyralid

RISK EXPRESSIONS RELATED TO APPLICATOR AND ENVIRONMENT:

R38: Irritant to skin.
R41: Serious damage risk on eye
R52/53: Toxic for aquatic organisms, may cause adverse effects longtime in aquatic environment

SAFETY EXPRESSIONS:

S2: Keep out of the reach of children
S25: Avoid contact with eye.
S26: In case of contact with eye, wash plenty of water immediately and call a doctor.
S29: Do not empty into drains
S37: Wear suitable gloves.
S39: Wear protective glass / mask
S46: Call a doctor immediately if swallowed and show the container or label.
S56: Dispose of this material and container in collection places for dangerous or special wastes.

MANUFACTURING AND PERMIT HOLDING COMPANY:
AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.
Bekir Saydam Cad. No: 31
Pancar, Torbali, Izmir
Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

(Company Stamp & Signature)

Time of Application:

Korazim should be applied after Beet and Creeping Thistle emerges. For best results application should be performed between 2-4 leaved stage of Creeping Thistle until it reaches 15-20 cm in height.

INFORMATION ON RESISTANCE:

The pesticide name **Korazim** is a herbicide classifies as Group O in accord with its action mechanism. Repeated applications of pesticides with same action mechanisms invite development of resistance. Therefore, to delay development of resistance, do not exceed the recommended number of **Korazim** applications in the same season. In case repeated application is necessary, take care to use pesticides with a different action mechanism (except of Group O).

MISCIBILITY:

Korazim can be applied in mixture with herbicides containing 2,4-D Amin, 2,4 D Isooctyl ester 2,4-D acid Isooctyl ester+trichlorpyr butoxyethylester, an MCPA.

Signature and Stamp
Dr. Yunus BAYRAM
Department Chief

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



KORAZIM

Soluble Concentrate (SL)

Name & Amount of Active Ingredient: 100 g
Clopyralid in Liter
Clopyralid (3.6-Dichloropicolinic acid) 3.6-
dichloropyridine-2-carboxylic acid

GROUP	O	HERBICIDE
--------------	----------	------------------

SYMPTOMS OF POISONING:

Headache, dizziness, weakness, nausea and haze

FIRST AID MEASURES:

Take the patient away from the contaminated area. Take off the contaminated clothes. In case of contact with skin, wash with plenty of water and soap. In case of contact with eye, rinse the eye with water for 15 minutes. The doctor can decide whether the pesticide is removed from the stomach.

ANTIDOTE AND NECESSARY INFORMATION:

There is no special antidote.

**NATIONAL POISON INFORMATION CENTRE (UZEM)
PHONE: 114**

5 December 2014
MINISTRY OF FOOD AGRICULTURE AND LIVESTOCK
GENERAL DIRECTORATE OF FOOD AND CONTROL
ACCEPTED

HANDLING AND STORAGE CONSIDERATIONS:

- Store the pesticide in its closed packaging.
- Suitably dispose the empty packages.
- Do not use empty packages for any other purpose.
- Do not put pesticide in empty food or drink containers.
- Toxic for bees.
- Do not use during florescence
- Toxic for fish
- Avoid contaminating above and ground waters.

STORAGE CONDITIONS:

In its original package store it closed , away from food and Keep locked up and out of the reach of children and in cool and dry place. When stored under cool and dry conditions in its original closed package, the product will not experience any physical, chemical or chemical changes outside the tolerance limits for 2 years.

DISCLAIMER:

The manufacturer warrants the quality of the product as long as it is sold in its original packaging. The user is responsible for the damage risk, arising from the factors out of the manufacturer. The manufacturer shall not accept any responsibility regarding damages originating from erroneous storage or failure to follow instructions.

MANUFACTURING AND PERMIT HOLDING COMPANY:
AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.
Bekir Saydam Cad. No: 31
Pancar, Torbali, Izmir
Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

(Company Stamp & Signature)