

1

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

# SPARTA 25 WP

Group:3;G1, Fungicide

Powder Wettable in Water

Active Ingredient:25% Tebuconazole (RS)-1-p-chlorophenyl-4, 4-dimethyl-3-(1H-1,2,4-triazol-1-ylmethyl) pentan-3-ol

**READ THE LABEL FIRST**

**DO NOT USE AT HOME**

**KEEP AWAY FROM CHILDREN, FOOD AND ANIMAL FOOD**

**DO NOT INHALE THE VAPOR AND DRIBLETS**

**WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES**

**DO NOT EAT, DRINK OR SMOKE DURING THE APPLICATION**

**AREA APPLIED IS NON-ACCESSIBLE FOR HUMANS AND ANIMALS FOR 1 DAY**

**FOLLOW THE USING INSTRUCTIONS IN ORDER TO PREVENT RISKS ON HUMAN AND ENVIRONMENT HEALTH**

**THE USE ON THE PRODUCTS OTHER THAN RECOMMENDED STRICTLY PROHIBITED**

**TOXICATION SYMPTOMS:**

Difficulty in transpiration and mobility, cramps.

**FIRST AID PRECAUTIONS:**

Call doctor, keep the patient away from the area of effect. The doctor must see the product package or label.

**ANTIDOTE:** No special antidote. Symptomatic treatment is followed.

NATIONAL TOXIC CONSULTANCY ASSOCIATION (NPCA) Tel:114

## 2

Production date:

Expiry date:

Charge nr:

License date and nr:

### PLANTS and DISEASES TO APPLY ON

Plant	Disease	Dosage and Timing	Period between final spraying and harvest		
apple	podospaera leucotricha	25 g/100 l water	14 days		
	venturia inaequalis	25 g/100 l water			
pear	venturia pirina	25 g/100 l water	14 days		
apricot	Sclerotinia laxa	60 g/100 l water	14 days		
Cereals	Puccinia striiformis	75 g/da			
	Septoria tritici	75 g/da	----		

tomato	Alternaria solani	50 g/100 l water	7 days		

**Preparing for applying:** First, make a mixture with a little amount of water in a separate tank. Fill in the half of the tool tank, and add the mixture into it, then add into the necessary amount of water while the mixer is on.

**Tomato:**By the time the primal symptoms are noticed, the plants are directly coating sprayed every 10-14 days.

**Pear:**1.spraying, by the time the flower buds bloom.

2.spraying at the rosette leaf term

3.spraying, by the time the 80% of the petals fall, and repeated every 10 days according to the weather conditions.

### 3

**Apricot:**1. spraying, at the beginning of blooming (on 5-10% flowers)

2. spraying, at full blooming (on 90-100% flowers)

**Cereals:**

First spraying as coating, by the time the primal symptoms are noticed around. Provided that the conditions are convenient to the disease, a second spraying is made after 10-14 days.

**Apple:**

**Against powdery mildew (*podosphaera leucotricha*):**

1. spraying at the pink rosette step
2. spraying, by the time the 60-70% of petals fall
3. spraying and the others each 10 days

**Against black spot (*venturia inaequalis*):**

1. spraying by the time the buds bloom.
2. spraying at pink rosette term, by the time the 70-80% falls.
3. spraying and the others, every 15 days provided that the weather conditions are convenient for the disease to develop.

#### INFORMATION ON IMMUNITY :

Sparta 25 WP is a fungicide classified as Group 3, G1 according to its effect mechanism. Repetitive applications of the plant protect products with the same effect mechanism encourages a development of immunity. In order to delay this development, do not exceed the total number of application amount advised for the season. In case a repeat is needed, use plant protect product with different effect mechanism (out of the Group 3, G1).

**MISCIBILITY:**Can be mixed with WP formulations. A pre-trial must be made with the other formulations. In general, mixing with highly acidic and basic formulations is not possible.

## 4

T:TOXIC  
N:DANGEROUS FOR THE ENVIRONMENT  
CONTAINS TEBUCONAZOLE

### RISK JARGON AND EXPLANATIONS

R22:Toxic if inhaled  
R63:Risk of harming the unborn infant.  
R51:Toxic for aquatic organisms  
R53:May result in long term negative effects in the aquatic environments.

### SAFETY JARGON AND EXPLANATIONS

S1/2:Keep under lock, out of the reach of children  
S9:Store the package at a very well aerated Place.  
S13:Keep away from food, beverage, and animal food  
S36/37:Wear appropriate protective uniform, and gloves.  
S46:If swallowed, immediately see a doctor, show the product package and label.  
S29:Do not empty into canalisation.  
S56:Have this product and its package exterminated at a waste treatment facility  
S61:Avoid release into the nature.Follow the using instructions/Safety Information Form

MANUFACTURER AND LICENSED COMPANY:  
AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.  
Bekir Saydam Cad. No:31,  
Pancar Beldesi, Torbalı-İZMİR  
Tel:0232 864 13 33, Fax:0232 864 16 79

### USAGE and STORAGE FACTS:

- Keep the product in its package, closed
- Exterminate accordingly the emptied packages
- Do not use the empty packages for any other purposes
- Do not refill the emptied food and beverage cans with the product.
- Toxic for fish.
- Avoid contaminating into underground and surface waters.

**STORING:**

Keep in the original package, closed, away from food, out of reach of children, and animals, at a cool and dry place, under lock. If stored under normal conditions (cool and dry), in original package, and closed keeps its physical, and chemical properties for min. 2 years within tolerance limits.

**COMPANY'S ANNOUNCEMENT:**

The company guarantees quality provided that the product is sold in original package.  
The company denies all the responsibility for the damages that may result of storing the product under wrong conditions or not following the recommendations while applying.

MANUFACTURER AND LICENSED COMPANY:  
AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.  
Bekir Saydam Cad. No:31,  
Pancar Beldesi, Torbalı-İZMİR  
Tel:0232 864 13 33, Fax:0232 864 16 79