

1

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

PAROMET 50 EC

Emulsion Concentrate (EC)

Active Ingredient:500 g Imazalil /L

(±)-1-(β-allyloxy-2,4-dichlorophenylethyl)imidazole; allyl (±)-1-(2,4-dichlorophenyl)-2-imidazol-1-ylethyl ether

GROUP; 3; G1; FUNGICIDE

CAUTION

READ THE LABEL FIRST.

DO NOT USE AT HOME.

KEEP AWAY FROM CHILDREN AND FOOD.

DO NOT INHALE THE VAPOR AND DRIBLETS

WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES

DO NOT EAT, DRINK, OR SMOKE DURING THE APPLICATION

AREA APPLIED IS NON-ACCESSIBLE FOR HUMANS AND ANIMALS FOR A DAY

FOLLOW THE USING INSTRUCTIONS IN ORDER TO AVOID RISKS ON HUMAN AND ENVIRONMENT HEALTH

THE USE ON THE PRODUCTS OTHER THAN RECOMMENDED STRICTLY PROHIBITED

TOXICATION SYMPTOMS:

Trembling time to time, dizziness, itch, red marks on skin and diarrhea. High loss of appetite, nausea, and vomiting.

FIRST AID PRECAUTIONS:

Consult a doctor and show him the product package or label.

If contacted skin; take off the contaminated uniforms immediately, wash well the skin surfaces that contacted with water and soap.

In case of eye contact; wash well with water, wahs well min. for 15 minutes also while the eyes are open. Consult a doctor immediately.

NATIONAL TOXIC CONSULTANCY ASSOCIATION (NPCA) Tel:114

2

Production date:

Expiry date:

Charge nr:

License date and nr:

PLANTS and DISEASES TO APPLY ON

Plant	Disease	Dosage and Timing	Period between final spraying and harvest		
grape	botrytis cinerea	30 ml/100 L water	3 days		
tomato	botrytis cinerea	30 ml/100 L water	3 days		
	Alternaria solani	30 ml/100 L water	3 days		

Preparing for applying: First, make a mixture with water in a separate tank. Fill in the half of the tool tank, and add the mixture into it, then add water into until it is completely full, while the mixer is on. In case of spraying, the plant must be washed in the best way. By the help of under foliar nozzles, the back surfaces must also be wetted. The sprayings must be made on days that are not windy.

WHEN TO USE

botrytis cinerea, grapes:The cure must be started right before the start of the ripening of the grains, and must be repeated every 10-15 days.

botrytis cinerea, tomato:The sprayings must be started by the time the primal symptoms are noticed, and they must be repeated every 10-15 days.

Alternaria solani, tomato: The sprayings must be started by the time the primal symptoms are noticed, and they must be repeated every 10 days.

3

IMMUNITY:

Paromet 50 EC is a fungicide classified as Group 3, G1 according to its effect mechanism. Repetitive applications of the plant protect products with the same effect mechanism encourages a development of immunity. In order to delay this development, do not exceed the total number of application amount of Paromet 50 EC advised for the season. In case a repeat is needed, use plant protect product with different effect mechanism (out of the Group 3, G1).

MISCIBILITY:

Can be mixed with Captan, Carbendazim compounds.

4

Net Amount:

Gross Amount:

Min. Retail Sales Price:

Xn:Hazardous

N:DANGEROUS FOR THE ENVIRONMENT

CONTAINS Imazalil, and Xylene

RISK JARGON AND EXPLANATIONS

R10:Flammable

R20/21/22:Hazardous on health if inhaled, contacted with skin, or swallowed.

R38:Irritates skin

R41:Risk of serious damage on eye

R50:Very toxic for the aquatic organisms.

R53:May result in long term negative effects in the aquatic environments.

SAFETY JARGON AND EXPLANATIONS

S1/2:Keep under lock, out of the reach of children

S13:Keep away from food, beverage, and animal food

S25:Avoid eye contact

S26:Wash well with water in case of eye contact, and immediately consult a doctor.

S29:Do not empty into canalisation.

S36/37/39:Wear appropriate protective uniform, and gloves, protective glasses/mask.

S46:If swallowed, immediately see a doctor, show the product package and label.

S56:Have this product and its package exterminated at a waste treatment facility

S64:If swallowed, rinse mouth well with water (If the patient is conscious)

P11: For use of Professional consumers.

5

USAGE and STORAGE FACTS:

- Toxic for fish. Do not mix into waters.
- Consider the periods that should be between the last spraying and harvest.
- Keep the product in its package, closed.
- Exterminate the emptied bottles accordingly.
- Do not use the empty packages for any other purposes.
- Do not refill the emptied food and beverage cans with the product.
- Toxic for fish.
- Avoid contaminating into the underground and surface waters.
- Protect from fire, spark or excessive heat.

STORING:

Keep in its original package, closed, away from food, out of the reach of children and animals, under lock. If stored under normal conditions (cool and dry), in original package, and closed, keeps its physical, and chemical properties for min. 2 years within tolerance limits.

COMPANY'S ANNOUNCEMENT:

The company guarantees quality provided that the product is sold in original package.

The damages may be caused by the factors out of the manufacturer's control are at the user's risk.

The company denies all the responsibility for the damages that may result of storing the product under wrong conditions or applying it at wrong doses and times.

MANUFACTURER AND LICENSED COMPANY:

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79