

TRANSLATION

Manufacture Date :
 Expiration Date :
 Charge No :
 Permit Date & Number :

TARGETED PLANTS AND PESTS

PLANT NAME	PEST NAME	DOSAGE	LAST APPLICATION-HARVEST INTERVAL
VINEYARD	Mildew (<i>Plasmopara viticola</i>)	175 g/ 100 L water	14 days
	Dead arm (<i>Phomopsis viticola</i>)		
APRICOT	Ringed spot (<i>Spiloceae oleagina</i>)	175 g/ 100 L water	14 days
CITRUS	Brown (<i>Alternaria solani</i>)	200 g/ 100 L water;	14 days
OLIVE	Ringed spot (<i>Spiloceae oleagina</i>)	175 g/ 100 L water	14 days
TOMATO	Mildew (<i>Phytophthora infestans</i>)	200 g/ 100 L water;	14 days

USAGE METHOD:

Preparation of Pesticide:

Fill the tank of device in half with water. Mix the required pesticide with water in other container. and then add this mixture to this tank. At this time mixer system should be operating state. Then full the tank with water. Do not mix cyclafon with dilution whose pH is less than 6,5. Otherwise they together can cause burning.

Application time:

Mildew on vineyard: Start to disinfest when offshoots are 25-30 cm, continue 7-10 days apart in accordance with air conditions and situation of illness. Stop disinfesting when proper situations don't exist for illness infection anymore. By disinfestation, pay attention to disinfest to leaf and bunches which are on the inner side. Disinfest particularly soffits of leaves. Disinfestations must be done on mornings at early hours when weather is not so hot.

Dr.NEVZAT BIRISIK
General Manager Deputy
STAMP AND SIGNATURE

Continued on Page 2

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



Suspension Concentrate (SC)

Name & Amount of Active Ingredient:

Copper Hydroxide equal to 361,1 g/L Metallic Copper

GROUP	M1	FUNGICIDE
-------	----	-----------

FIRSTLY READ THE LABEL.
 DO NOT USE AT HOME.
 KEEP AWAY FROM CHILDREN, FOODSTUFFS AND ANIMAL FEED.
 DO NOT INHALE VAPORS AND/OR PARTICLES OF PESTICIDE.
 PLEASE USE PROTECTIVE CLOTHING, GLOVES AND GLASSES.
 DO NOT EAT, DRINK OR SMOKE DURING APPLICATION OF PESTICIDE.
 NO LIMITATION REQUIRED FOR HUMAN AND ANIMAL ENTRY INTO PESTICIDE TREATED AREAS FOR 2 DAYS.
 PLEASE FOLLOW THE USAGE INSTRUCTIONS TO PREVENT ANY HUMAN HEALTH AND ENVIRONMENTAL RISKS.
 IT IS STRICTLY PROHIBITED TO USE THE PESTICIDE ON ANYTHING OTHER THAN RECOMMENDED TARGETS.
 AFTER ADDING CLEAN WATER TO EMPTY CONTAINER OF PESTICIDE WITH THE RATE OF 1/4 AND SHAKE WELL. POUR THE MIXED SHAKEN WATER INTO DISINFESTION TANK. REPEAT THIS PROCESS THREE TIMES

NATIONAL TOXIN HOTLINE (UZEM) PHONE: 114

MINISTRY OF FOOD, AGRICULTURE AND LIVESTOCK
 GENERAL DIRECTORATE OF FOOD AND CONTROL
 ACCEPTED
 12 APRIL 2013

Net Amount :
 Max. Retail Sales Price :

HAZARD TYPES:

N: DANGEROUS FOR THE ENVIRONMENT

HAZARD SYMBOLS:



N: DANGEROUS FOR THE ENVIRONMENT

INGREDIENTS:

CONTAINS COPPER HYDROXIDE

RISK EXPRESSIONS:

R50/53: Plenty of toxic for aquatic organism can cause adverse effect in aquatic environment for a long time.

SAFETY EXPRESSIONS:

S1/2: Keep locked up and out of the reach of children
S9: Keep the tank in an environment which is ventilated very well.
S29: Do not empty into sewage.
S56: Dispose of this material and its container at hazardous or special waste collection point

HANDLING AND STORAGE CONSIDERATIONS:

- Store the pesticide in its closed packaging.
- Suitably dispose the empty packages.
- Do not use empty packages for any other purpose.
- Do not put pesticide in empty food or drink containers.
- Toxic for fish
- Avoid contaminating above- and underground waters.

MANUFACTURING AND PERMIT HOLDING COMPANY:
 AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.

Bekir Saydam Cad. No: 31
 Pancar, Torbali, Izmir, TURKEY
 Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

(Company Stamp & Signature)

Dead Arm in Vine yard: Summer disinfection is applied.

1st disinfection is applied when the twigs are at 2-3 cm of length.

2nd disinfection is applied when the twigs are at 8-10 cm of length.

3rd disinfection is applied when the twigs are at 25-30 cm of length.

Leaf piercing in apricot:

1st disinfection is applied after the defoliation in autumn (Dormant period)

2nd disinfection is applied before the flower buds are opened. (Normal period)

3rd disinfection is applied when the epicalyces and anther structures are pulled away. (Normal period)

Brown stain disease in citrus: The application starts when the first disease symptoms take places within the twigs in spring. It's repeated in ach ten days.

Mildew in tomatoes: The application starts when the symptoms take places. It's applied for twice in ach ten days.

INFORMATION ON RESISTANCE:

The pesticide name **Lampris** is a fungicide classified as Group M1 in accord with its action mechanism. Repeated applications of pesticides with same action mechanisms invite development of resistance. Therefore, to delay development of resistance, do not exceed the recommended number of **Lampris** applications in the same season. In case repeated application is necessary, take care to use pesticides with a different action mechanism (other than Group M1).

MISCIBILITY:

It can be used by being mixed with many pests excluding Carbaryl featured pests. Pre-mixture trial in farmer conditions must be performed just in case.

12 APRIL 2013

Dr.NEVZAT BIRISIK
General Manager Deputy
STAMP AND SIGNATURE

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



Water dissolvable dry fluid granule (DF)

Name & Amount of Active Ingredient:

Copper Hydroxide %53,8 (equal to %35 metallic copper)

GROUP	M1	FUNGICIDE
-------	----	-----------

SYMPTOMS OF POISONING:

Metallic taste in the mouth, bloody diarrhea, dizziness, weakness, blurred vision, anaemia.

FIRST AID MEASURES AND ANTIDOTE:

Call a doctor if swallowed. Make the patient vomit by making him/her drink 1-2 glasses of milk. Do not vomit him/her if the patient is unconscious. Wash the skin contacted parts with plenty of water and soap. Wash with plenty of water in case contact with eyes. Bring the doctor the pesticide package and label.

ANTIDOTE AND IMPORTANT INFORMATION:

There is no special antidote.

NATIONAL TOXIN HOTLINE (UZEM) PHONE: 114

MINISTRY OF FOOD, AGRICULTURE AND LIVESTOCK
GENERAL DIRECTORATE OF FOOD AND CONTROL
ACCEPTED
12 APRIL 2013

STORAGE CONDITIONS:

In its original package store it closed, away from food and keep locked up and out of the reach of children and in cool and dry place. When stored under normal (cool and dry) conditions in its original closed package, the product will not experience any physical, chemical or chemical changes outside the tolerance limits for 2 years.

DISCLAIMER:

The manufacturer warrants the quality of the product as long as it is sold in its original packaging. User endures damage risk which can occur because of factors out of manufacturer control. The manufacturer shall not accept any responsibility regarding damages originating from erroneous storage or failure to follow instructions.

MANUFACTURING AND PERMIT HOLDING COMPANY:

AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.
Bekir Saydam Cad. No: 31
Pancar, Torbali, Izmir, TURKEY
Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

(Company Stamp & Signature)

I hereby certify that this document has been translated from its Turkish into English truly and correctly by me.
İşbu belge Türkçe aslından İngilizce'ye tarafımdan aslına uygun olarak tercüme edilmiştir. 14.08.2013