

Production Date :  
Expire Date :  
Batch Number :  
License Date and Number. :

**PLANTS AND PESTS TO BE USED FOR**

Plant	PEST	USAGE DOSAGES	WAITING PERIOD
Tomato	early blight ( <i>Alternaria solani</i> )	200 g/ 100 l water	14 days
	Bacterial Pinta ( <i>Pseudomonas syringae</i> pv. <i>tomato</i> )	200 g/ 100 l water	14 days
Vineyard	Mildew ( <i>Plasmopara viticola</i> )	150 g / 100 l water	21 days
	dead arm ( <i>Phomopsis viticola</i> )	150 g/ 100 l water	21 days
APPLE	Black spot ( <i>Venturia inaequalis</i> )	200 g/ 100 l water	21 days

**USAGE:**

The recommended amount of medicine is slurried with little amount of water in a different cup, after than it is poured into the disinfection reservuar which is half-filled. Mixer of the device should be worked during the reservuar is filled fully with water.

For tomato, against early blight disinfection is started when the first indications of the disease appeared. Disinfection is continued with 10 days interval.

For tomato against ice-minus bacteria green assembly disinfections, when the first indications of diseas appeared for prevention, once a week for seed phase and with 8-10 days interval for land phase 2-3 application should be done.

Grape Yard mildew: Disinfection is started when offshoots are 25-30 cm long Subsequent disinfections are continued with 10-14 days intervals. According to the situation of the disease disinfections are continued until 21 days before crop.

Grape Yard antraknoz: 1. Summer disinfection when the offshores are 2-3 cm, 2. Summer disinfections when offshores are 8-10 cm and 3. Summer disinfection when the ofshores are 25-30 cm, should be applied.

Apple Blackspot Bruising: 1. Disinfection when the flower cells are swelled, 2. Disinfection when pink flower colliculus period

3. Disinfection when 70 – 80 % of the flower petals are fallen, 4. and other disinfections are continued with 10-15 days intervals in the occasions when the ecological conditions are proper for the progression of disease.

**AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.**

# EFFICURE 75 WG



Water Soluble Granul (WG)

**Name and quantity of the active component: % 75 Mancozeb**

Manganese ethylenbis (dithiocarbamate)polymeric  
mixture with zinc salt

GROUP	M3	FUNGICIDE
-------	----	-----------

**READ THE LABEL FIRST.**

**DO NOT USE AT HOME.**

**KEEP OUT OF REACH OF CHILDREN, FOOD AND ANIMAL FEED.**

**AVOID İNHALATION OF VAPORS OR PARTICULES.**

**WEAR MASK, PROTECTIVE CLOTH, GLOVES, GOGGLES.**

**DO NOT EAT, DRINK OR SMOKE DURING TREATMENT.**

**DO NOT USE THE MEDICATED SEEDS AS FOOD OR ANIMAL**

**FEED.**

**OBAY INSTRUCTIONS FOR USE TO AVOID THE RISK OF HUMAN AND ENVIRONMENTAL HEALTH.**

**USING TOGETHER WITH THE PRODUCTS NOT ADVISED**

**CERTAINLY FORBIDDEN.**

**TOXICOLOGICAL SYMPHOMS:** nausea, vomiting, dizziness, headache, skin mucous membrane irritation, fatigue, and blurred vision.

**FIRST AID MEASURES:** In case poisoning call a doctor, and shom the label. If contact the skin, take off dirty clothes and wash exposed skin with plenty of soapy water. In case contact of eye, wash with plenty of water for 15 minutes. In case of swallowed, have drunken 2 bottles of water.

**ANTIDOTE AND REQUIRED INFORMATION:** No specific antidote is known. Treatment symptomatically.

**NATIONAL POISON INFORMATION OFFICE: 114**


Net Weight :

Maximum Selling Price( inc. VAT):

**HAZARD PROPERTY**

Xn:Harmful N: Harmful for the Environment

**DANGER SYMBOLS**



Xn:Harmful

N: Harmful for the Environment

**INGREDIENTS:**

**It contains mancozeb**

**RISK PHRASES AND DISCLOSURES**

**R37** Irritating to respiratory system

**R43** May cause sensitization by skin contact.

**R63:** Possible risk of harm to the unborn child.

**R50** Very toxic to aquatic organisms.

**SAFETY PHRASES AND DISCLOSURES**

**S1/2:** Keep out of reach of children

**S9** Keep container in well-ventilated place.

**S13:** Keep away from.Food, drink and animal feedstuffs

**S20** When using do not eat or drink.

**S24** Avoid contact with skin.

**S29** Do not empty into drains.

**S36/37/39:** Use suitable protective clothing, gloves and protective goggles/mask.

**S46:** If swallowed, go to doctor immediately and show this container or label.

**S56:**Remove this material and its jar in special waste collection point.

**S63** If swallow, rinse mouth with water (Only if the person is conscious).

**PRODUCER AND OWNER OF THE LICENCE:**

**AGRI SCIENCES Tarım ve İlaç. Ür.**

San. ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31 Pancar Torbalı, İZMİR

Tel: 0090 0232 864 13 33

Fax: 0090 0232 864 16 79

**INFORMATION ABOUT RESISTANCE:**

**Effucure 75 WG** named plant protection products according to its mechanism of action is classified as Group M3. Repetitive applications of plant protection products with the same mechanism of action, encourages the development of resistance. For this reason, to delay development of resistance in **Effucure 75 WG** on the same production season, the number of applications exceeds the recommended total. In cases where the application should be repeated, with a different mechanism of action (Group M3 external) careful use of plant protection products.

**MIXING CASES: Effucure 75 WG** mixes with many fungicides, insecticide foliar fertilizers and plant growth regulators. See statements of mixed before mixture. **Effucure 75 WG** mixes active matters contents drug such as captan, carbendazim, dinocop, folpet, cymoxanil and festyl al.

**AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.**



# EFFICURE 75 WG



Water Soluble Granul (WG)

**Name and quantity of the active component:** % 75 Mancozeb

Manganese ethylenbis (dithiocarbamate)polymeric)  
mixture with zinc salt

GROUP	M3	FUNGICIDE
-------	----	-----------

**NATIONAL POISON INFORMATION OFFICE: 114**

**SAFETY PRECAUTIONS DURING USING AND STORAGE:**

- \* Keep in closed original container
- \* Dispose of empty containers properly.
- \* Do not use empty containers for other purposes.
- \* Do not put food and drink boxes, empty plant protection product.
- \* Harmful for bees
- \* Do not use on blooming period
- \* Harmful for fishes
- \* Avoid infecting ground and surface water.

**STORAGE CONDITIONS:**

Store in dry and cool conditions, in closed original container separately from feed and food, If stored in cool and dry conditions, the product will keep its physical, chemical and biological structure at least 2 years.

**COMPANY DECLARATION:**

The quality of the pesticide is guaranteed by the producer if it is to be sold in the original packaging. The producer will not be responsible of the losses and damages resulting from the handling and storage mistakes or wrong treatment or improper actions not mentioned on the label.

**PRODUCER AND OWNER OF THE LICENCE:**

**AGRI SCIENCES Tarım ve İlaç Ür.**

San. ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31 Pancar Torbalı, İZMİR

Tel: 0090 0232 864 13 33

Fax: 0090 0232 864 16 79