

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

TRINITY

Herbicide

Emulsion Concentrated

Active ingredient:330 g Pendimethanil / liter

N-(1-ethylpropyl)-2,6-dinitro-3,4-xylidine

Acute oral in rats LD50 < 1340 mg/kg

TOXIC

READ THE LABEL FIRST.

DO NOT USE AT HOME.

KEEP AWAY FROM CHILDREN AND FOOD.

AVOID SKIN AND EYE CONTACT

WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES

DO NOT EAT DURING THE APPLICATION, DO NOT SMOKE

DO NOT MAKE SPRAYING AGAINST WIND

DO NOT INHALE THE VAPOR AND DRIBLETS

WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES

DO NOT EAT DURING THE APPLICATION

AFTER SPRAYING, WASH WELL THA AREAS THAT CONTACTED

THE PRODUCT WITH WATER AND SOAP

TOXICATION SYMPTOMS:

Irritation in eyes and on skin, engorgement, fever.

FIRST AID PRECAUTIONS:

If swallowed, do not make the patient vomit. Call a doctor immediatley. In case of eye or skin contact, wash well with water min. for 15 minutes and call the doctor.

ANTIDOTE:

No special antidote. Symptomatic treatment is followed.

TREATMENT:

If swallowed, immediately maket he patient plenty of milk, egg white or if these are inavailable, water. Avoid alcohol. Call the doctor for medical treatment.

NATIONAL TOXIC CONSULTANCY ASSOCIATION (NPCA) Tel:114

Production date:
Expiry date:
Charge nr:
License date and nr:

PLANTS TO APPLY ON and DISEASES

PLANT	SAVAGE HERBS	DOSE
SUNFLOWER, COTTON, ONION, POTATO	Large and small leaf herbs Alopecurus myosuroides Digitaria sanguinalis Poa pratensis Amaranthus spp. Atriplex spp. Chenopodium album Euphorbia spp. Heliotropium europaeum Polygonum spp. Solanum nigrum Sisymbrium officinale Urtica urens Papaver rhoeas Apera spica-venti Echinochloa crus-galli Setaria spp. Anagallis arvensis Capsella bursa-pastoris Chrosphora tinctoria Setyaria verticillata Lamium purpureum Portulaca oleracea Stellaria media Thlaspi arvense Matricaria spp. Sorghum halepense (from seed)	500 ml/da
MAIZE, TOMATO	Large and small leaf herbs Amaranthus spp. Chenopodium album Solanum nigrum Echinochloa colonom Portulaca oleracea Thlaspi arvense Chrosphora tinctoria	500 ml/da

***Do not use for plants that grow in very sandy soil .**

GENERAL INFORMATION:TRINITY is a herbicide that controls the large and small leaf herbs of diversives of cultivated plants. It can be mixed into the soil befor the planting, can be used before or after the emergence.

TRINITY, controls the savage herbs in case of sprouting or right after it.

APPLICATION:TRINITY must be applied by ground tools with 20-50 l water per decar, and if possible, shall be sprayed by fan type nozzles with 110°.

3 (page 2 continues)

Preparing:

- 1.Fill in the half of the tank with water. If there exits, turn on the mixer of the tool.
- 2.Add the necessary amount of TRINITY into the tank.
- 3.Fill in the rest of the rest of the tank with water.
- 4.Mix well, then start spraying.

HOW TO APPLY:

1.FOR COTTON AND SUNFLOWER:Applying TRINITY by mixing it into soil gives the best results. Spraying can be done until 7 days before the planting. The field must be clod free and well prepared. Within 4 days that follow the spraying, mixing into soil must be until 5-8 cm of depth, and by disc, rototiller, or harrow.If there happens a rainfall of 10 mm within 4 days after the spraying, mixing into soil is not necessary. For applying onto cotton, must be applied within 5 days after the planting, before the emergence.

2.FOR POTATO, MAIZE, ONION AND TOMATO GROWN FROM SEED:TRINITY is applied onto the soil surface after the seed planting at pre-emergence. The depth of planting the seeds must be 4-5 cm deep. For the potatoes, it must be applied at the term before the emergence of the potatoes and the savage herbs. For the maize, TRINITY is never mixed into soil and it is never used for sweet corn. Before using on the maize fields that have been planted for the seed reproduction, please contact the AGRI SCIENCES competents. For the onions, can be at pre-emergence, and also at double leaf step as well, but provided that it is before the herbs pass to the 1-2 foil step. The tomato seeds must be planted between the 2. and 6. days after the spraying. In case of planting the seeds, the contact of the roots with the soil that has been mixed with the product must be avoided.

UNDER DRY CLIMATE CONDITIONS, 5-10 mm WATERING MUST BE DONE WITHIN 3-7 DAYS THAT FOLLOW THE APPLYING OF TRINITY.

MISCIBILITY:If it is intended to make tank mixture of TRINITY with other products, its physical miscibility must be controlled.

ROTATION PLANTS:After a spraying with TRINITY, no rotation happens for 6 months with any plant except the following. Cotton, peas, maize, soybean, peanuts, sunflower, and dilution plants. When the cultivated plant goes off with any kind of reason, potato, beans, summer cereals, maize, cotton, soybean, peas can be planted into min. 15 cm deep. Maize, beans and peas can be planted before the soil is treated, provided that the depth is min. 5 cm.

4

Net amount:

Gross amount:

Max. Retail sales price (VAT included):

Date and nr of the Ministry of Agriculture's Letter

Shake well before use.

USAGE AND STORAGE FACTS:

1. Seedbed must be well prepared, the soil must be fine and clod free.
2. The advised planting depths must be strictly followed.
3. Although TRINITY is much effective than many other soil affected herbicide, some amount of humidity is necessary to provide the full effect.
4. When used for soils that contain organic materials more than 10%, there can be a decrease in control of the herbs.
5. Must not be used at surface pools.
6. TRINITY must not be used for the plants that are weak and stressed resulting of diseases, drought, excessive chemical application, insufficient soil preparation, etc.
7. TRINITY cannot control the herbs of many hears.
8. The liquid and the vapour of TRINITY is flammable.
9. Toxic for the fish, must not be contaminated into waters.
10. The emptied packages must be exterminated accordingly, the tool must be washed well.

LICENSED COMPANY:

Gennova Tarım İlaçları Sanayi ve Ticaret AŞ.

Cumhuriyet Bul. No:280/101

Alsancak/İZMİR

Tel:0232 464 23 85, Fax:0232 464 23 95

MANUFACTURER COMPANY:

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79

5

STORING:

If stored under normal conditions (cool and dry), in the original package, keeps its physical, chemical, and biologic properties of for min. 2 years within the tolerance limits.

COMPANY'S ANNOUNCEMENT:

The instructions on the label concerning the usage of this product reflect the related researches made by experts and field application results. The correctness and reliability of the instructions have been priorly approved, and the using advices are must does. On the other side, eliminating all the risks that may occur related to the use of the product is impossible. These may be factors such as weather conditions, existence of other products or usage mistakes whih are out of the control of AGRI SCIENCES, that may end in damage, inefficiency and undesired results. Such risks are to the user. AGRI SCIENCES only guarantees the conformity of the mentioned product to the chemical specifications mentioned on the label and the product is conforming to the purpose of use except the above mentioned risks, provided that the using instructions are followed.

LICENSED COMPANY:

Gennova Tarım İlaçları Sanayi ve Ticaret AŞ.

Cumhuriyet Bul. No:280/101

Alsancak/İZMİR

Tel:0232 464 23 85, Fax:0232 464 23 95

MANUFACTURER COMPANY:

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79