

1

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

PAVLINA 100 EC

Fungicide

Emulsion Concentrated

Active Ingredient and ratio:contains 100 g/l **PENCONAZOLE**

1-(2-(2,4-dichlorophenyl) pentyl)- 1 H – 1,2,4-triazole

LD₅₀ : 2574 mg/kg

TOXIC

READ THE LABEL FIRST

DO NOT USE AT HOME

KEEP OUT OF REACH OF CHILDREN, ANIMAL, AND ANIMAL FOOD

DO NOT INHALE THE PRODUCT VAPOR AND ITS DRIBLET

AVOID SKIN AND EYE CONTACT

WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES

DO NOT EAT DURING THE APPLICATION, DO NOT SMOKE

HUMANS AND ANIMALS FOR ONE DAY ARE NOT ALLOWED INTO THE AREA APPLIED

THE USE ON THE PRODUCTS OTHER THAN RECOMMENDED STRICTLY PROHIBITED

TOXICATION SYMPTOMS:

No toxication on human has been recorded so far.

FIRST AID PRECAUTIONS:

Despite it is not considered as possible, if toxication is doubted, call a doctor. Stop patient's contact with the product. Take off all the uniforms, and wash well with water and soap. Rinse the eyes min. for 15 minutes with water. Take the patient to a well aerated place.

ANTIDOTE :No any specific antidotes. Treatment according to the symptoms must be followed.

NPCA (NATIONAL TOXIC CONSULTANCY ASSOCIATION) Tel:114

2

Production date:

Expiry date:

Charge nr:

License date and nr:

THE PLANTS AND DISEASES TO APPLY ON

Plant	Disease	Application Dose and Term	Period Between Last Spraying and Harvest(days)
Orchard	Unciuia necator	25 ml/100 l water or decar	21
Roses	Spnaerochea pannosa	25 ml/100 l water	21

or decar

WHEN TO USE:

Spraying are started before the flowering, by the time the first leaves develop and the buds reach to a length of 20-30 cm. This way, it is prevented that the disease factor reaches to a high level. 4-5 spraying are recommended during the season.

The intervals between spraying are advised as 14 days for the orchard and the roses.

PREPARING TO APPLY:

First the nozzle of the tool is controlled and cleaned well, so that it runs properly. Then, the half of the tank is filled with water. The necessary amount of the product is mixed with a little amount of water in a separate tank and added into. Then more water is added the tank is completely full, while the mixer of the apparel is on. For a good coating, the calibration of the tool must have been fixed. The spraying must be done when it is not windy.

MISCIBILITY

PAVLINA is not affected by the other product applied before or after itself, as well as it is not affected of those with which it is applied as a tank mixture. If needed, a pre-mixture is advised of those that will be applied as mixture.

PAVLINA is a fungicide that is absorbed rapidly into the metabolism right after it is applied, and stored in the leaves. After it is pulverised onto the plants, it is not washed away by the rains.

3

Net Amount:

Gross Amount:

Max. Retail Sales Price:

The date and the number of the Letter of the Ministry of Agriculture and Rural Affairs:

CAUTIONS ON USAGE AND STORAGE

*Toxic for the fish.

*Animals should not be pastured on, crops (neither for the human nor the animals) shouldn't be collected from, and the pets should be kept away from the newly sprayed field.

*Keep the product in the original package, closed, away from the food, out of reach of children.

*Do not use the emptied product bottles for any other purpose and exterminate accordingly.

STORING:

Provided that the product is kept in the original package, closed and at a dry and cold place, keep its physical, chemical, and biologic properties for two years.

*Keep the products that are not used tightly close, out of reach of children, away from food, under lock.

*Toxic for bees. Do not use at flowering time.

*Toxic for the fish, do not contaminate into the waters.

*Must be stores at cool and dry places.

*Exterminate the emptied bottles accordingly.

COMPANY'S ANNOUNCEMENT:

The company guarantees the quality of the product provided that it is sold in the original and closed package, but denies the responsibility for the damages that may result of storing or applying the product in a wrong way or not following the advices properly.

LICENSED COMPANY:

Genova Tarım İlaçları Sanayi ve Ticaret AŞ.
Cumhuriyet Bul. No:280/101
Alsancak/İZMİR
Tel:0232 464 23 85, Fax:0232 464 23 95

MANUFACTURER COMPANY:

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.
Bekir Saydam Cad. No:31,
Pancar Beldesi, Torbalı-İZMİR
Tel:0232 864 13 33, Fax:0232 864 16 79