

Production Date :
Expire Date :
Batch Number :
License Date and Number. :

PLANTS AND PESTS TO BE USED FOR

Plant	Pest	Usage dosages	The period between the last spray and harvest
Vineyard	Grape powdery mildew (<i>Uncinula necator</i>)	400 g / 100 l water	7 days
Fruit	Apple Powdery Mildew (<i>Podosphaera leucotricha</i>)	400 g / 100 l water	7 days
	Peach Powdery Mildew (<i>Sphaerotheca pannosa</i>)	400 g / 100 l water	7 days
Ornamental Plants	Rose Powdery Mildew (<i>Sphaerotheca pannosa</i>)	400 g/ 100 l water	-----
	Begonia Powdery Mildew (<i>Oidium Begoniae</i>)	300-350 g/ 100 l water	-----
Vegetable	Cucurbitaceae Powdery Mildew (<i>Erysiphe cichoracearum</i>)	400 g/100 l water	7 days
	Chickpea Anthracnose (<i>Ascochyta rabiei</i>)	300 g/ 100 l water	7 days
	Solanaceae Powdery Mildew (<i>Leveillula taurica</i>)	400 g/100 l water	7 days
	Onions Downy Mildew (<i>Peronospora destructor</i>)	300 g / 100 l water	7 days
	Bean Rust (<i>Uromyces appendiculatus</i>)	300 g/100 l water	7 days
VINEYARD	Grape Erineum Mite (<i>Eriophyes vitis</i>)	400 g/ 100 l water	7 days
FRUIT TREES	Fruit Tree's Mites (<i>Tetranychus viennensis</i> , <i>T.urticae</i>), <i>Panonychus ulmi</i>)	400 g/100 l water	7 days
CITRUS	Citrus red Mite (<i>Panonychus citri</i>)	600 g/ 100 l water	7 days
	Citrus Bud Mite (<i>Aceria sheldoni</i>)	600 g/100 l water	7 days

USAGE INSTRUCTIONS

Preparing the medicine for use:

Sufficient amount of medicine is mixed with some water in a container. Half of the medicine apparatus is filled with water and rest of it is filled with the medicine mix prepared.

Time of Use:

Grape Yard powdery mildew:

- 1.Disinfection when offshoots are 25-30 cm long
- 2.Disinfection when grapes are small
- 3.Disinfection depending on the progress of the disease and meteorological and ecological conditions of the environment. When the grape pieces candied.

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

AGRI SCIENCES
DIAFLEX

Water Dispersible Granule (WG)

Name and quantity of the active component: %80 Sulfur
Micronized Sulfur

GROUP	M2	FUNGICIDE - ACARICIDE
-------	----	-----------------------

READ THE LABEL FIRST.
DO NOT USE AT HOME.
KEEP OUT OF REACH OF CHILDREN, FOOD AND ANIMAL FEED. AVOID INHALATION OF VAPORS OR PARTICULES. WEAR MASK, PROTECTIVE CLOTH, GLOVES, GOGGLES. DO NOT EAT, DRINK OR SMOKE DURING TREATMENT. DO NOT USE THE MEDICATED SEEDS AS FOOD OR ANIMAL FEED.
OBEY INSTRUCTIONS FOR USE TO AVOID THE RISK OF HUMAN AND ENVIRONMENTAL HEALTH.
USING TOGETHER WITH THE PRODUCTS NOT ADVISED CERTAINLY FORBIDDEN.
NATIONAL POISON INFORMATION OFFICE: 114

TOXICOLOGICAL SYMPHOMS: Chilling result of influencing of central nervous system, in further lethargy or convulsions.
FIRST AID MEASURES: In case contact the skin, take off dirty clothes and wash exposed skin with plenty of soapy water. In case contact of eye, wash with plenty of water for 15 minutes. In case of swallowed, call a doctor, show the label or the package.
ANTIDOTE AND REQUIRED INFORMATION: No specific antidote is known.

NATIONAL POISON INFORMATION OFFICE: 114

Net Weight :
Maximum Selling Price:

HAZARD PROPERTY

Xi: Irritant

DANGER SYMBOLS



Xi: Irritant

INGREDIENTS:

It contains sulfur

RISK PHRASES AND DISCLOSURES

R38 Irritating the skin.

SAFETY PHRASES AND DISCLOSURES

S1/2: Keep out of reach of children
S13: Keep away from.Food, drink and animal feedstuffs
S36/37/39: Use suitable protective clothing, gloves and protective goggles/mask.
S46: If swallowed, g oto doctor immediately and show this container or label.
S64 If swallow, rinse mouth with water (Only if the person is conscious).

PRODUCER AND OWNER OF THE LICENCE:

AGRI SCIENCES Tarım ve İlaç. Ür.

San. ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31 Pancar Torbalı, İZMİR

Tel: 0090 0232 864 13 33

Fax: 0090 0232 864 16 79

Apple powdery mildew:

1. Disinfection when pink flower colliculus period
2. Disinfection when 60 – 70 % of the flower petals are fallen
3. Other disinfections: once per week till the end of May, once per 10 days till 3rd week of July.

Peach powdery mildew:

1. Disinfection period must be decided according to intensity of the disease at the previous years. If no previous disease, disinfection is made when first indication of the disease appears. Other infections: according to growth of the plant and progress of the disease, intervals of 8-12 days till the disease is controlled.

Rose powdery mildew: 1st disinfection is made when leaves and collicula first appears. According to the progress of the disease, 5-6 times of disinfection is made.

Begonia powdery mildew: When the indications of the disease first appear, once per week

Cucurbitaceae Powdery Mildew: The first symptoms of the disease seem in the region, starts to spraying. According to the course of the disease, sparing With an interval of 10 days.

Chickpea antracnose: When the indications of the disease first appear, once per 7-10 days depending on the intensity of the disease and meteorological conditions for 2-5 times.

Eggplant powdery mildew: When the indications of the disease first appear, once per 10 days depending on the intensity of the disease and meteorological conditions for 3-5 times.

Onion powdery mildew: When the indications of the disease first appear, once per 10 days depending on the intensity of the disease and meteorological conditions for 3-5 times.

Bean Rust: When the indications of the disease first appear, once per week for 3-5 times.

INFORMATION ABOUT RESISTANCE:

Diaflex named plant protection products according to its mechanism of action is classified as Group M2. Repetitive applications of plant protection products with the same mechanism of action, encourages the development of resistance. For this reason, to delay development of resistance in **Diaflex** on the same production season, the number of applications exceeds the recommended total. In cases where the application should be repeated, with a different mechanism of action (Group M2 external) careful use of plant protection products.

MIXING CASES: It is not mixed with wintery and summary oil. A germination test should be performed first, if the mixture.

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.



Water Dispersible Granule (WG)

Name and quantity of the active component: %80 Sulfur
Micronized Sulfur

GROUP	M2	FUNGUCIDE - ACARICIDES
-------	----	---------------------------

NATIONAL POISON INFORMATION OFFICE: 114

SAFETY PRECAUTIONS DURING USING AND STORAGE:

- * Keep in closed original container
- * Dispose of empty containers properly.
- * Do not use empty containers for other purposes.
- * Do not put food and drink boxes, empty plant protection product.
- * it doesn't harmful for bees and fishes
- * Cucurbitaceous and apricots sulfur is sensitive.
- * It should be after at least 1 month from applied.

STORAGE CONDITIONS:

Store in dry and cool conditions, in closed original container separately from feed and food, If stored in cool and dry conditions, the product will keep its physical, chemical and biological structure at least 2 years.

COMPANY DECLARATION:

The quality of the pesticide is guaranteed by the producer if it is to be sold in the original packaging. The producer will not be responsible of the losses and damages resulting from the handling and storage mistakes or wrong treatment or improper actions not mentioned on the label.

PRODUCER AND OWNER OF THE LICENCE:

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.
Bekir Saydam Cad. No:31 Pancar Torbalı, İZMİR
Tel: 0090 0232 864 13 33
Fax: 0090 0232 864 16 79