

Production Date:
Expire Date:
Batch No:
License Date and No:

PLANTS AND PESTS TO BE USED FOR

Plant	Pest	Usage dosages	The period between the last spray and harvest
Apple	Codling moth (<i>Cydia Pomenella</i>)	20 ml/ 100 l water	14 days
	Webworm (<i>Yponomeuta Malinellus</i>)	20 ml/100 l water	14 days
	* <i>Stigmella malella</i> (<i>Leucoptera scitella</i>)	20 ml/100 l water	14 days
Pear	Pear leaf psyllid (<i>Cacopsylla pyri</i>)	25 ml+50ml spreader for 100 l water	14 days
Cotton	Cotton Leaf worm (<i>Spodoptera littoralis</i>)	30 ml/deccare	14 days

(*) There is no necessary to apply insecticide for also against *Stigmella malella* in gardens fight against Codling moth.

PREPARING THE PRODUCT :

Fill the half of the tank of the application device with clean water. Mix necessary amount of pesticide in another clean container with few water. Then pour that solution into the tank of the application device while its agitator is on. Add necessary amount of water while still mixing. Spraying should be done windless weather.

AGRI SCIENCES TARIM VE İLAÇ ÜR. SAN. VE TİC. LTD. ŞTİ.



AFFLURON

Suspension Concentrate (SC)

Name and quantity of the active component:480 g/L
Diflubenzuron 1-(4-chlorophenyl)-3-(2,6-difluorobenzoyl)urea

GROUP	15	INSECTICIDE
-------	----	-------------

READ THE LABEL FIRST.

DO NOT USE AT HOME.

KEEP OUT OF REACH OF CHILDREN, FOOD AND ANIMAL FEED. AVOID İNHALATION OF VAPORS OR PARTICULES.

WEAR MASK, PROTECTIVE CLOTH, GLOVES, GOGGLES. DO NOT EAT, DRINK OR SMOKE DURING TREATMENT. DO NOT ALLOW HUMAN AND ANIMAL FOR 1 DAY TO ENTER TO THE AREA WHERE APPLICATION DRUG.

OBEY INSTRUCTIONS FOR USE TO AVOID THE RISK OF HUMAN AND ENVIRONMENTAL HEALTH. USING TOGETHER WITH THE PRODUCTS NOT ADVISED CERTAINLY FORBIDDEN.

TOXICOLOGICAL SYMPHTOMS:

Headache, dizziness and vomiting

FIRST AID MEASURES

Wash the contacting parts of body with water and soup. If it contact to eye, wash with water for 15 minutes, call medical assistance and show the label or the package.

ANTIDOTE: No specific antidote is known. symptomatic treatment.

NATIONAL POISON INFORMATION OFFİCE: 114

Net Weight :
Maximum Selling Price(inc. VAT):

HAZARD PROPERTY

N: Harmful for the Environment

DANGER SYMBOLS



N: Harmful for the Environment

INGREDIENTS:

It contains diflubenzuron.

RISK PHRASES AND DISCLOSURES

R50 Very toxic to aquatic organisms.

SAFETY PHRASES AND DISCLOSURES

S1/2: Keep out of reach of children

S13: Keep away from.Food, drink and animal feedstuffs

S29 Do not empty into drains.

S56:Remove this material and its jar in special waste collection point.

S61 Avoid release to the environment. Refer to special instructions and safety data sheets.

PRODUCER AND OWNER OF THE LICENCE:

AGRI SCIENCES Tarım ve İlaç. Ür.

San. ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31 Pancar Torbalı, İZMİR

Tel: 0090 0232 864 13 33

Fax: 0090 0232 864 16 79

APPLICATION TIME:

Apple codling moth control : Control and counts in sexual traps are taken into account for on every trust to start spraying. Starting to spraying in according to butterfly flights, damages, biology, phenological state of the trees, and the information received from organizations.

Apple webworm control: A single spraying is done when larvae began to weave first networks after leave leaf epidermis.

Stigmella malella control: Controls that can be started from the beginning of the month of April, almost all of the eggs left off wintering adults of reproductive age 2, and 3 in a single Pesticide application when it starts seeing nymphs sufficient.

Cotton Leaf worm control: When shown 0.5 larvae per plant or 2 eggs for 25 plants or 2 new dehiscence, dressing begins.

INFORMATION ABOUT RESISTANCE:

Affluron named plant protection products according to its mechanism of action is classified as Group 15. Repetitive applications of plant protection products with the same mechanism of action, encourages the development of resistance. For this reason, to delay development of resistance in Affluron on the same production season, the number of applications exceeds the recommended total. In cases where the application should be repeated, with a different mechanism of action (Group 15 external) careful use of plant protection products.

MIXING CASES

Generally, it can be used with other pesticides that have the form of same formulation. Attempts to pre-mixture is recommended, before large amounts of mixture for spraying.

AGRI SCIENCES TARIM VE İLAÇ ÜR. SAN. VE TİC. LTD. ŞTİ.



AFFLURON

Suspension Concentrate (SC)

Name and quantity of the active component:480 g/L
Diflubenzuron 1-(4-chlorophenyl)-3-(2,6-difluorobenzoyl)urea

GROUP	15	INSECTICIDE
--------------	-----------	--------------------

NATIONAL POISON INFORMATION OFFİCE: 114

SAFETY PRECAUTIONS DURING HANDLING AND STORAGE

- * Keep in closed original container
- * Dispose of empty containers properly.
- * Do not use empty containers for other purposes.
- * Do not put food and drink boxes, empty plant protection product.
- * Harmful for fishes
- * Avoid infecting ground and surface water.

STORAGE CONDITIONS:

Keep it in original closed packaging, away from food, out of reach of children, cool and dry place. If stored in cool and dry conditions, the product will keep its physical, chemical and biological structure at least 2 years.

COMPANY DECLARATION:

The quality of the pesticide is guaranteed by the producer if it is to be sold in the original packaging. The producer will not be responsible of the losses and damages resulting from the handling and storage mistakes or wrong treatment or improper actions not mentioned on the label.

PRODUCER AND OWNER OF THE LICENCE:

AGRI SCIENCES Tarım ve İlaç. Ür.

San. ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31 Pancar Torbalı, İZMİR

Tel: 0090 0232 864 13 33

Fax: 0090 0232 864 16 79