

Manufacture Date :  
 Expiration Date :  
 Charge No :  
 Permit Date & Number :

**TARGETED PLANTS AND PESTS**

PLANT NAME	PEST NAME	DOSAGE & STAGE	LAST APPLICATION-HARVEST INTERVAL
VINEYARD	Botrytis cinerea	75 g/100 L water	14 days
TOMATO	Botrytis cinerea	75 g/100 L water	7 days
	Early blight (Alternaria solani)	100 g/100 L water	
CUCUMBER	Sclerotinia sclerotiorum	75 g/100 L water	7 days
APRICOT	Monillia laxa	150 g/100 L water	14 days
CITRUS	Alternaria alternate f.sp.citri	100 g/100 L water	14 days

**USAGE METHOD:**

**Preparation of Pesticide:**

Fill the tank of device in half with water. Melt required pesticide in other container. Operate mixer of tool and add pesticide and full it with water. Use prepared solution within the same day. Disinfestation must be done on a windless weather. Pay attention to get every side of plant wet but pesticide shouldn't flow. Every side of bunches in vineyard must be disinfested.

**Application Time:**

**Botrytis cinerea in vineyard:** Disinfestation should be started just before beginning of maturing of grapes.

**Botrytis cinerea in tomato:** Disinfestation must be started when first symptoms are seen or when plants are on blossoming period, the second disinfestation is done after 10 days.

**Early blight in tomato:** It is started when first stains are seen.

**Dr.NEVZAT BIRISIK**  
**General Manager Deputy**  
**STAMP AND SIGNATURE**

Continued on Page 2

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



Wettable powder (WP)

**Name & Amount of Active Ingredient:**

% 50 Iprodione  
 3-(3,5- dichlorophenyl)- N- isopropyl-2,4- dioxoimidazolidine-1-carboxamide

GROUP	2, E3	FUNGICIDE
-------	-------	-----------

FIRSTLY READ THE LABEL.  
 DO NOT USE AT HOME.  
 KEEP AWAY FROM CHILDREN, FOODSTUFFS AND ANIMAL FEED.  
 DO NOT INHALE VAPORS AND/OR PARTICLES OF PESTICIDE.  
 PLEASE USE PROTECTIVE CLOTHING, GLOVES AND GLASSES.  
 DO NOT EAT, DRINK OR SMOKE DURING APPLICATION OF PESTICIDE.  
 NO LIMITATION REQUIRED FOR HUMAN AND ANIMAL ENTRY INTO PESTICIDE TREATED AREAS FOR 4 HOURS.  
 PLEASE FOLLOW THE USAGE INSTRUCTIONS TO PREVENT ANY HUMAN HEALTH AND ENVIRONMENTAL RISKS.  
 IT IS STRICTLY PROHIBITED TO USE THE PESTICIDE ON ANYTHING OTHER THAN RECOMMENDED TARGETS.

**NATIONAL TOXIN HOTLINE (UZEM) PHONE: 114**

MINISTRY OF FOOD, AGRICULTURE AND LIVESTOCK  
 GENERAL DIRECTORATE OF FOOD AND CONTROL  
 ACCEPTED  
 19 DECEMBER 2012

Net Amount :  
 Max. Retail Sales Price :

**HAZARD TYPES:**

Xn: Harmful (carcinogenic, cat.3) N: DANGEROUS FOR THE ENVIRONMENT

**HAZARD SYMBOLS:**



Xn: Harmful



N: DANGEROUS FOR THE ENVIRONMENT

**INGREDIENTS:**

CONTAINS IPRDIONE

**RISK EXPRESSIONS:**

**R40:** Limited evidence for carcinogenic effect.

**R50/53:** Toxic for aquatic organisms, it can cause adverse effects longtime in aquatic environment.

**SAFETY EXPRESSIONS:**

**S1/2:** Keep locked up and out of the reach of children

**S13:** Keep away from food, drink and animal feeding stuffs

**S29/56:** Do not empty into drains. Dispose of this material and its container at hazardous or special waste collection point

**S36/37:** Wear suitable protective clothing, gloves and eye/face protection.

**S46:** Call a doctor if swallowed and show the container or label.

**MANUFACTURING AND PERMIT HOLDING COMPANY:**  
**AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.**

Bekir Saydam Cad. No: 31

Pancar, Torbali, Izmir, TURKEY

Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

(Company Stamp & Signature)

**Sclerotinia sclerotiorum in cucumber:** Disinfestation is started when first stains are seen.

**Monillia laxa in apricot:**

1. Disinfestation is done at the beginning of blossoming (%5-10)

2. Disinfestation is done (%90-100) at perfect flower.

**Alternaria alternate f.sp.citri in citrus: Disinfestation is done when meeting is seen 10 days apart.**

**INFORMATION ON RESISTANCE:**

The pesticide name **Hickory** is an insecticide classifies as Group 2, E3 in accord with its action mechanism. Repeated applications of pesticides with same action mechanisms invite development of resistance. Therefore, to delay development of resistance, do not exceed the recommended number of **Hickory** applications in the same season. In case repeated application is necessary, take care to use pesticides with a different action mechanism (other than Group 2, E3).

**MISCIBILITY:**

It can be used by being mixed with fungicides, insecticides and foliar fertilizers except Bordeaux mixture. Before doing huge amount of mixture, it is advised to make physical miscibility test.

**Dr.NEVZAT BIRISIK**  
**General Manager Deputy**  
**STAMP AND SIGNATURE**

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



Wettable powder (WP)

**Name & Amount of Active Ingredient:**

%50 Iprodione

3-(3,5- dichlorophenyl)- N- isopropyl-2,4- dioximidazolidine-1-  
carboxamide

GROUP	2, E3	FUNGICIDE
-------	-------	-----------

**SYMPTOMS OF POISONING:**

Tonus, reflex decay and imbalances in manners.

**FIRST AID MEASURES AND ANTIDOTE:**

In case of swallow, call a doctor, puke the patient or wash the stomach. In case of contact with skin, wash with plenty of water and soap. Take off the contaminated clothes. In case contact with eye, wash with plenty of water and apply an eye doctor. if the patient is conscious, take the patient in lateral position and carry.

**ANTIDOTE AND IMPORTANT INFORMATION:**

There is no special antidote. In accord with symptoms supportive treatment applied.

**NATIONAL TOXIN HOTLINE (UZEM) PHONE: 114**

MINISTRY OF FOOD, AGRICULTURE AND LIVESTOCK  
GENERAL DIRECTORATE OF FOOD AND CONTROL  
ACCEPTED  
19 DECEMBER 2012

**HANDLING AND STORAGE CONSIDERATIONS:**

- Store the pesticide in its closed packaging.
- Suitably dispose the empty packages.
- Do not use empty packages for any other purpose.
- Do not put pesticide in empty food or drink containers.
- Avoid contaminating above- and underground waters.

**STORAGE CONDITIONS:**

In its original package store it closed , away from food and dry place. When stored under normal (cool and dry) conditions in its original closed package, the product will not experience any physical, chemical or chemical changes outside the tolerance limits for 2 years.

**DISCLAIMER:**

The manufacturer warrants the quality of the product as long as it is sold in its original packaging. The manufacturer shall not accept any responsibility regarding damages originating from erroneous storage or failure to follow instructions.

**MANUFACTURING AND PERMIT HOLDING COMPANY:**

**AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.**  
Bekir Saydam Cad. No: 31  
Pancar, Torbali, Izmir, TURKEY  
Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

(Company Stamp & Signature)