

Manufacture Date :  
 Expiration Date :  
 Charge No :  
 Permit Date & Number :

**TARGETED PLANTS AND PURPOSE, PERIOD AND DOSAGE**

PLANT NAME	APPLICATION METHOD & PERIOD	DOSAGE
<b>Seedless grape for table consumption</b>	It is used to enlarge bunches and grapes. Used when flower petals are defoliated 70%	100 ml/100 L water
	Used after 10-12 days after the first application when the grapes reach to 3 mm diameter	100 ml/100 L water
<b>Seedless grape for raisin production</b>	Used a couple of days after the flower petals are defoliated	25 ml/100 L water
<b>CITRUSES</b> TANGERINE	Especially used on Satsuma and Clementine species of tangerine at 75-80% blossom stage to stimulate pollination.	75 ml / 100 L water
ORANGE	Used 4-5 weeks after full blossom on orange species excluding Valencia.	100 ml / 100 L water
WASHINGTON ORANGE	Used on Washington orange when color fruit starts to change to delay the harvest.	50 ml / 100 L water
LATE SEASON LEMON	To delay harvest: Used when fruit color first starts to change from green to yellow, in order to ensure fruit can stay on branch by 2-3 months more.	50 ml / 100 L water
GRAPEFRUIT	To delay harvest: Used when fruit color first starts to change from green to yellow, in order to ensure fruit can stay on branch by 2-3 months more.	75 ml / 100 L water
<b>PEAR</b>	Used at 30% blossom to increase yield in pear species with low pollination rate and low fruit development rate. Also used in 2 days after frost at the latest to protect against frost damages that can occur during flowering period.	75-100 ml / 100 L water
<b>PEACH</b>	An application of <b>Geno-Gibb Max</b> solution in first week of August, especially on Dixired and Hale species, prevents the peach from being damaged by late spring frosts and automatically provides thinning.	50-75 ml / 100 L water
<b>STRAWBERRY</b>	Used 3-4 weeks before blossoming to increase fruit size and yield.	75-100 ml / 100 L water
<b>ARTICHOKE</b>	Used especially on heart of the plant when leaf length is 25-30 cm to obtain early artichoke and to ensure large and regular shaped flower.	125-150 ml / 100 L water
<b>CHERRY</b>	Used when color fruit starts, in order to change to obtain durable, large and glossy cherries.	75-100 ml / 100 L water
<b>POTATO</b>	Potato seed stock is immersed in <b>Geno-Gibb Max</b> solution before planting to ensure quick and regular emergence. This expedites germination and regular emergence.	25 ml / 100 L water
<b>LETTUCE &amp; CURLY LETTUCE</b>	Used after seedlings are transplanted to field to increase yield.	125-250 ml / 100 L water
<b>OTHER PRODUCE</b>	<b>Geno-Gibb Max</b> can also be used on many other plants like ornamental plants, long pepper, lettuce, melon, watermelon, parsley, cucumber, squash, banana, eggplant, tomato, pea, bean, etc.	-

**THE WAITING PERIOD BETWEEN LAST APPLICATION AND HARVEST IS "ZERO" DAYS. ATTENTION: IT IS ESPECIALLY RECOMMENDED TO USE SPREADER ADHESIVE IN GENO-GIBB MAX APPLICATIONS.**

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



# GENO-GIBB MAX

Soluble Concentrate (SL)

**Name & Amount of Active Ingredient:**  
 16.66 g Gibberellic acid (GA<sub>3</sub>) in Liter  
 (3S, 3aR,4S,4aS,7S,9aR,12S)-7,12 dihydroxy-3methyl-6-methylene-2-oxoperhydro-4a,7-methano-9b,3-propenoazuleno(1,2-1,2-b)furan-4-carboxylic acid

<b>GROUP</b>	----	<b>PGD</b>
--------------	------	------------

FIRSTLY READ THE LABEL.  
 DO NOT USE AT HOME.  
 KEEP AWAY FROM CHILDREN, FOODSTUFFS AND ANIMAL FEED.  
 DO NOT INHALE VAPORS AND/OR PARTICLES OF PESTICIDE.  
 PLEASE USE PROTECTIVE CLOTHING, GLOVES AND GOGGLES.  
 DO NOT EAT, DRINK OR SMOKE DURING APPLICATION OF PESTICIDE.  
 DO NOT ALLOW HUMAN AND ANIMAL TO ENTRY INTO PESTICIDE TREATED AREAS FOR 1 DAY  
 PLEASE FOLLOW THE USAGE INSTRUCTIONS TO PREVENT ANY HUMAN HEALTH AND ENVIRONMENTAL RISKS.  
**IT IS STRICTLY PROHIBITED TO USE THE PESTICIDE ON ANYTHING OTHER THAN RECOMMENDED TARGETS.**  
 SHAKE WELL AFTER ADDING CLEAN WATER TO EMPTY CONTAINER OF PESTICIDE WITH THE RATE OF 1/4. POUR THE SHAKEN WATER INTO DISINFECTION TANK. REPEAT THIS PROCESS THREE TIMES

NATIONAL POISONING INFORMATION CENTRE (UZEM) : 114

Net Amount :  
 Max. Retail Sales Price :

**HAZARD TYPE**  
 Xn: Harmful

**HAZARD SYMBOLS AND EXPLANATION:**



Xn: HARMFUL

**INGREDIENTS:**  
 Gibberellic acid and diethylene glycol

**RISK EXPRESSIONS:**  
 R22: Harmful if swallowed.

**SAFETY EXPRESSIONS:**  
 S1/2: Keep locked and out of the reach of children  
 S13: Keep away from food, drink and animal feeding stuffs.  
 S36/37/39: Wear suitable clothes, protective clothes and protective glass/mask  
 S46: Call a doctor immediately if swallowed and show the container or label.

**HANDLING AND STORAGE CONSIDERATIONS:**

- Store the pesticide in its closed packaging.
- Suitably dispose the empty packages.
- Do not use empty packages for any other purpose.
- Do not put pesticide in empty food or drink containers.

**MANUFACTURING AND PERMIT HOLDING COMPANY:**  
 AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.  
 Bekir Saydam Cad. No: 31  
 Pancar, Torbali, Izmir  
 Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

**METHOD OF APPLICATION:**

**Preparation:** Dissolve recommended dosage of **Geno-Gib Max** in a small amount of water in a separate container. Pour this mixture into the applicator tank and fill the tank with water. Add necessary amount of spreader adhesive. Use the mixture without delay.

**INFORMATION ON RESISTANCE:**

The pesticide name **Geno-Gibb Max** is a Plant Growth Designator classifies as Group -- in accord with its action mechanism. Repeated applications of pesticides with same action mechanisms invite development of resistance. Therefore, to delay development of resistance, do not exceed the recommended number of **Geno-Gibb Max** applications in the same season. In case repeated application is necessary, take care to use pesticides with a different action mechanism (except of Group --).

**MISCIBILITY:** It can be used by mixing with fungicides, insecticides and leaf fertilizers except of the solutions including alkali features and chlorines

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.

**GENO-GIBB MAX**

Soluble Concentrate (SL)

**Name & Amount of Active Ingredient:**

Litrede 16.66 g Giberellic asit (GA<sub>3</sub>) in Liter  
(3S, 3aR,4S,4aS,7S,9aR,12S)-7,12 dihydroxy-3methyl-6-methylene-2-oxoperhydro-4a,7-methano-9b,3-propenoazuleno(1,2-1,2-b)furan-4-carboxylic acid

<b>GROUP</b>	----	<b>PGD</b>
--------------	------	------------

**SYMPTOMS OF POISONING:**

There is no special poisoning symptoms.

**FIRST AID MEASURES:**

Wash with plenty of water if contaminating eye and body.

**ANTIDOTE AND NECESSARY INFORMATION:**

There is no special antidote. The treatment is performed as per the symptoms.

NATIONAL POISONING INFORMATION CENTRE (UZEM) : 114

**STORAGE CONDITIONS:**

In its original package store it closed, away from food and keep locked up and out of the reach of children and in cool and dry place. When stored under cool and dry conditions in its original closed package, the product will not experience any physical, chemical or chemical changes outside the tolerance limits for 2 years.

**DISCLAIMER:**

The manufacturer warrants the quality of the product as long as it is sold in its original packaging. The user is responsible for the damage risk, arising from the factors out of the manufacturer. The manufacturer shall not accept any responsibility regarding damages originating from erroneous storage or failure to follow instructions.

**MANUFACTURING AND PERMIT HOLDING COMPANY:  
AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.**

Bekir Saydam Cad. No: 31

Pancar, Torbali, Izmir

Tel: +90-232-864-13-33 Fax: +90-232-864-16-79