

Manufacture Date :
 Expiration Date :
 Charge No :
 Permit Date & Number :

TARGETED PLANTS AND PESTS

PLANT NAME	PEST NAME	DOSAGE & TIME PERIOD	TIME BETWEEN FINAL APPLICATION & HARVEST
APPLE	Apple aphid (<i>Aphis pomi</i>)	40 ml/100 L water nymph & adult	14 days
	Codling moth (<i>Cydia pomonella</i>)	40 ml/100 L water Larva	
PEAR	Pear psyllid (<i>Cacopsylla pyri</i>)	50 ml/100 L water nymph	14 days
	Codling moth (<i>Cydia pomonella</i>)	40 ml/100 L water nymph	
QUINCE	Codling moth (<i>Cydia pomonella</i>)	40 ml/100 L water nymph	
CHERRY	Cherry fly (<i>Rhagoletis cerasi</i>)	40 ml/100 L water Adult	14 days
	European Leaf Roller (<i>Archips rosanus</i>)	50 ml/100 L water Larva	
PISTACHIO	Pistachio psyllid (<i>Agonosceca pistaciae</i>)	50 ml/100 L water nymph & adult	21 days
	Flat-headed sapling borer (<i>Capnodis cariose</i>)	50 ml/100 L water Adult	

METHOD OF APPLICATION:

Preparation:

First mix the necessary amount of product with water in a separate container. Fill the tank of applicator with water to the half point and pour the mixture in. Add the rest of necessary amount of water and continue mixing. Application should be performed in cool and calm weather, taking care ensure full coverage.

Time of Application:

Cherry fly control application should be performed in 7-10 days following the date when first adults are captured in visual yellow sticky traps.

Codling moth control applications should be coordinated according to forecast and warning system.

Apple aphid control application should be performed when 15% of the saplings are found to be infested.

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



Oil Based waterspension Concentrate (OD)

Name & Amount of Active Ingredient: 240 g

Thiacloprid in Liter

N-(3-(6-Chloro-pyridin-3-ylmethyl)-thiazolidin-2-ylidene)-cyanamide

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

FIRSTLY READ THE LABEL.

DO NOT USE AT HOME.

KEEP AWAY FROM CHILDREN, FOODSTUFFS AND ANIMAL FEED.

DO NOT INHALE VAPORS AND/OR PARTICLES OF PESTICIDE.

PLEASE USE PROTECTIVE CLOTHING, GLOVES AND GOGGLES.

DO NOT EAT, DRINK OR SMOKE DURING APPLICATION OF PESTICIDE.

DO NOT ALLOW HUMAN AND ANIMAL TO ENTRY INTO PESTICIDE TREATED AREAS FOR 1 DAY

PLEASE FOLLOW THE USAGE INSTRUCTIONS TO PREVENT ANY HUMAN HEALTH AND ENVIRONMENTAL RISKS.

IT IS STRICTLY PROHIBITED TO USE THE PESTICIDE ON ANYTHING OTHER THAN RECOMMENDED TARGETS.

SHAKE WELL AFTER ADDING CLEAN WATER TO EMPTY CONTAINER OF PESTICIDE WITH THE RATE OF 1/4. POUR THE SHAKEN WATER INTO DISINFECTION TANK. REPEAT THIS PROCESS THREE TIMES

NATIONAL TOXIN HOTLINE (UZEM) PHONE: 114

Net Amount :

Max. Retail Sales Price :

HAZARD TYPES:

Xn: Harmful N: DANGEROUS FOR THE ENVIRONMENT

HAZARD SYMBOLS



Xn: HARMFUL



N: DANGEROUS FOR THE ENVIRONMENT

INGREDIENTS:

Thiacloprid

RISK EXPRESSIONS RELATED TO APPLICATOR AND ENVIRONMENT:

R20/22: Harmful if inhaled or swallowed.

R40: Limited evidence for cancerogenic effect.

R51/53: Toxic for aquatic organisms, may cause adverse effects longtime in aquatic environment

SAFETY EXPRESSIONS:

S1/2: Keep locked and out of the reach of children

S9: Store the container in a well-ventilated area.

S13: Keep away from food, drink and animal feeding stuffs.

S29: Do not empty into drains

S35: Dispose of the material and container safely.

S37: Wear suitable clothes, protective gloves.

S46: Call a doctor immediately if swallowed and show the container or label.

MANUFACTURING AND PERMIT HOLDING COMPANY:

AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.

Bekir Saydam Cad. No: 31

Pancar, Torbali, Izmir

Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

Pear psyllid control application should be performed when 15% of the saplings are found to be infested and when 2nd and 3rd stage nymphs are in the majority.

Leaf roller application should be performed once when larva infestation rate in flower bunches reach 5% and ¾ of petals are shed.

Pistachio: Leaf psyllid control application should be performed when an average of 20-30 nymphs per leaf are counted.

Flat-headed sapling borer control application should be performed when adult population increases and when nibbles are found on fresh leaves at stem level or graft eyes or shoots.

INFORMATION ON RESISTANCE:

The pesticide name **Tackle** is an insecticide classifies as Group in accord with its action mechanism. Repeated applications of pesticides with same action mechanisms invite development of resistance. Therefore, to delay development of resistance, do not exceed the recommended number of **Tackle** applications in the same season. In case repeated application is necessary, take care to use pesticides with a different action mechanism (except of Group).

MISCIBILITY:

It can be mixed with many conventional pesticides, having formulation of EC, SL, WP and WG. However, a mixture test must be performed to be watered.



Oil Based waterspension Concentrate (OD)

Name & Amount of Active Ingredient: 240 g
Thiacloprid in Liter
N-(3-(6-Chloro-pyridin-3-ylmethyl)-thiazolidin-2-ylidene)-cyanamide

GROUP	INSECTICIDE
-------	-------------

SYMPTOMS OF POISONING:

Insensibility, spasm, hard to breath, trembling and cramp.

FIRST AID MEASURES:

Take the patient away from the contaminated area. Keep the patient horizontally in case of risk for faint. Take off the contaminated clothes. If inhaled, take the patient to the fresh air. In case of contact with skin, wash with plenty of water and soap. In case of contact with eye, rinse the eye with water well. If swallowed, do not induce vomiting. Rinse the mouth with water. Call a doctor or take the package or label of the pesticide with you.

ANTIDOTE AND NECESSARY INFORMATION:

There is no special antidote. The treatment is performed as per the symptoms.

NATIONAL TOXIN HOTLINE (UZEM) PHONE: 114

HANDLING AND STORAGE CONSIDERATIONS:

- Store the pesticide in its closed packaging.
- Suitably dispose the empty packages.
- Do not use empty packages for any other purpose.
- Do not put pesticide in empty food or drink containers.
- Toxic for bees.
- Do not use during florescence
- Toxic for fish
- Avoid contaminating above and underground waters.

CAUTION: Shake well before usage

STORAGE CONDITIONS:

In its original package store it closed , away from food and Keep locked up and out of the reach of children and in cool and dry place. When stored under cool and dry conditions in its original closed package, the product will not experience any physical, chemical or chemical changes outside the tolerance limits for 2 years.

DISCLAIMER

The manufacturer warrants the quality of the product as long as it is sold in its original packaging. The user is responsible for the damage risk, arising from the factors out of the manufacturer. The manufacturer shall not accept any responsibility regarding damages originating from erroneous storage or failure to follow instructions.

MANUFACTURING AND PERMIT HOLDING COMPANY:

AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.
Bekir Saydam Cad. No: 31
Pancar, Torbali, Izmir
Tel: +90-232-864-13-33 Fax: +90-232-864-16-79