

1

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

# VISTRO 500 SC

## Suspension Concentrate (SL)

Active Ingredient:500 g/L Clofentezine  
3,6-bis(2-chlorophenyl)-1,2,4,5-tetrazine

Propyl 3-(dimethylamino) propylcabamate hydrochloride

**GROUP: 10; A; ACARICIDE**

### CAUTION

READ THE LABEL FIRST.

DO NOT USE AT HOME.

KEEP AWAY FROM CHILDREN, FOOD, AND ANIMAL FOOD.

DO NOT INHALE THE VAPOR AND DRIBLETS.

WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES.

DO NOT EAT, DRINK, OR SMOKE DURING THE APPLICATION.

AREA APPLIED IS ACCESSIBLE FOR HUMANS AND ANIMALS.

FOLLOW THE USING INSTRUCTIONS IN ORDER TO AVOID RISKS ON HUMAN AND ENVIRONMENT HEALTH  
THE USE ON THE PRODUCTS OTHER THAN RECOMMENDED STRICTLY PROHIBITED

### TOXICATION SYMPTOMS:

No specific symptoms. Generally fatigue. When taken in high amounts, may cause eye and skin irritation.

### FIRST AID PRECAUTIONS:

Low toxicity for the human and animals, follow the general protection preventions. Wash well with soap and water if contacted skin. In case of toxication, rest the patient and immediately call a doctor, keep the patient away from the sprayed area. Shall the doctor see the product label and package.

### ANTIDOTE AND TREATMENT:

No specific antidotes. Symptomatic treatment is followed. Do not try to make the patient vomit. Give him gastronomic lavage with warm water.

NATIONAL TOXIC CONSULTANCY ASSOCIATION (NPCA) Tel:114

## 2

Production date:

Expiry date:

Charge nr:

License date and nr:

PLANTS and HAZARDOUS ORGANISMSTO APPLY ON

<b>Plant</b>	<b>Disease</b>	<b>Dosage and Timing</b>	<b>Period between final spraying and harvest</b>
apple	European red mite Red spider mite Hawthorn red spider mite	20 ml/100 L water	28 days
peach	Red spider mite	40 ml/100 L water	21 days
grape	Red spider mite	25 ml/100 L water	56 days
vegetable	Red spider mite Carmine spider mite	30 ml/100 L water	3 days
cotton	Red spider mite	40 ml/da	
soybean	Carmine spider mite	40 ml/da	

**HOW TO USE:**

Use once a year.

**On European red mite eggs:** Before the cracking of winter eggs, at the beginning of the pink bud term of the trees, spraying must be done. If this period is missed, spraying must be made when 1-2 mature larvae are seen per leaf.

**In gardens where Red spider mite, Hawthorn red spider mite, and others dominate:** Spraying must be done during the period starting with the fall of petals and ending with yeasting time.

**For Red spider mite eggs on grapes, vegetables, cotton:** Spraying must be done when the primal actions are noticed (1-2 mature larvae per leaf)

Fill the half of the tank with water. Then add the necessary amount of the acaricide, then add water while mixing.

In order to get the best results, a good coating must be done as if all the sides of the trees are being washed.

**IMMUNITY:**

Vistro 500 SC is a fungicide classified as Group 10, A according to its effect mechanism. Repetitive applications of the plant protect products with the same effect mechanism encourages a development of immunity. In order to delay this development, do not exceed the total number of application amount of Vistro 500 SC advised for the season. In case a repeat is needed, use plant protect product with different effect mechanism (out of the Group 10, A).

**MISCIBILITY:**

Can be used with other insecticides and fungicides.

**3****Net Amount:****Min. Retail Sales Price:****DANGER:**

N:DANGEROUS FOR ENVIRONMENT

Contains Clofentezine

**RISK JARGON AND EXPLANATIONS**

R52: Hazardous for the aquatic organisms.

R53: May result in long term negative effects in the aquatic environments.

## **SAFETY JARGON AND EXPLANATIONS**

S1/2:Keep under lock, out of the reach of children

S13:Keep away from food, beverage, and animal food

S29:Do not empty into canalisation.

S36/37/39:Wear appropriate protective uniform, and gloves, protective glasses/mask.

S56:Have this product and its package exterminated at a waste treatment facility

S64:If swallowed, rinse mouth well with water (If the patient is conscious)

## **USAGE and STORAGE FACTS:**

- Keep the product in its package, closed.
- Exterminate the emptied bottles accordingly.
- Do not use the emptied packages for any other purposes.
- Do not refill the emptied food and beverage cans with the product.
- Toxic for the fish.
- Avoid contaminating into the underground and surface waters.

## **STORING:**

Keep in its original package, closed, away from food, out of the reach of children, at a cool and dry place under lock. If stored under normal conditions (cool and dry), in original package, and closed, keeps its physical, and chemical properties for min. 2 years after the production date.

## **COMPANY'S ANNOUNCEMENT:**

The company guarantees quality provided that the product is sold in original package.

The damages may be caused by the factors out of the manufacturer's control are at the user's risk.

The company denies all the responsibility for the damages that may result of storing the product under wrong conditions or applying it at wrong doses and times.

## **MANUFACTURER AND LICENSED COMPANY:**

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79