

1

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

# CASPERTIN

**Wetable Powder**

**Group:17-Fungicide**

Effective Ingredient:50% Cymoxanil

Chemical:1-(2-cyano-2-methoxyiminoacetyl)-3-ethylurea

**READ THE LABEL FIRST. DO NOT USE AT HOME**

**KEEP AWAY FROM CHILDREN AND FOOD**

**DO NOT INHALE THE VAPOR AND DRIBLETS**

**WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES**

**AVOID SKIN AND EYE CONTACT .**

**DO NOT EAT DURING THE APPLICATION**

**DO NOT SMOKE**

**HUMANS and ANIMALS FOR ARE NOT ALLOWED INTO THE AREA APPLIED FOR 7 DAYS**

**FOLLOW THE INSTRUCTIONS OF USE IN ORDER TO PREVENTS RISKS ON HUMAN and ENVIRONMENT HEALTH**

**STRICTLY PROHIBITED TO USE ON CROPS OTHER THAN THE RECOMMENDED**

## **TOXICATION SYMPTOMS**

Headache,dizziness, fatigue, transpiration.

**FIRST AID PRECAUTIONS:**In case of toxication, take the following first aid preventions, call the doctor, let him see the product's package and labels. Wash well the surfaces that contacted the product with water and soap. Keep the patient away from the spraying area. Make him vomit if swallowed.

**ANTIDOTE:**No any special antidote.

NATIONAL TOXIC CONSULTANCY ASSOCIATION (NPCA) Tel:114

## 2

Production date:

Expiry date:

Charge nr:

License date and nr:

### PLANTS and INSECTS TO APPLY ON

Plant	Savage Herb	Dose	Period between last spraying and harvest
Zucchini	False dawning mildew ( <i>Pseudoperonospora cubensis</i> )	50 g/100 L water	3 days

### USING DIRECTIONS

#### Preparation to Apply

The spraying tank is filled in with water in half. In a separate tank, the product is mixed with some amount of water, then added into the spraying tank. It is mixed with mechanic or hidrolic mixers, and the rest of the water is also added into.

In case of spraying, the plant must be washed away in the best way, both the upper and bottom surfaces of the leaves must be wetted very well. Spraying must be done in cool and smooth weather conditions. The spraying for the dawning mildew in zucchini is started when the branches begin to grow or the first signs of the dawning mildew are observed. Min. two sprayings are done between periods of five days.

### INFORMATION ON RESISTANCE

Repetitive use of products having the same effect mechanism encourages a development of resistance. For this reason, to delay the resistance development, do not exceed the advised total number of applications of Caspertin within a single production period. In cases that requires a repeat of application, it is better to use the plant protect products with different effect mechanism (except Group 27)

### MISCIBILITY FACTS

Can be mixed with many other insecticides except the alkali such as high acid and lime or bordeaux mixes. But, before making a mixture of big amounts, the possibility of a physical mixture must be tested with a small amount of it.

### 3

Net Amount:  
Max. Sales price :

#### FACTS OF DANGER:

Xn:HAZARDOUS

N:DANGEROUS FOR NATURE

#### DANGER SYMBOL

.....  
Xn:HAZARDOUS

.....  
N:DANGEROUS FOR NATURE

#### INGREDIENTS:

Contains Cymoxanil.

#### JARGON TO RISK

R22:Hazardous if swallowed  
R43:May cause irritation if contacted with skin.  
R50:Very toxic for water organisms  
R53:May cause negative effects of long term at water .

#### JARGON TO SECURITY

S ½:Keep locked away fro children  
S22:Do not inhale the powder of it  
S24:Avoid contact with skin  
S36/37:Use appropriate protective uniform and gloves  
S46:See the doctor immediately if swallowed, show the package or label  
S29:Do not empty into the canalisation  
S56:Have this product and its package asided at a private or hazardous waste treatment facility.  
S61:Avoid releasing it into environment. Address to the special use instruction or the Security Information Form.

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.  
Bekir Saydam Cad. No:31,  
Pancar Beldesi, Torbalı-İZMİR  
Tel:0232 864 13 33, Fax:0232 864 16 79

### 4

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

**CASPERTIN**

**Wetable Powder****Group:17-Fungicide**

Effective Ingredient:50% Cymoxanil

Chemical:1-(2-cyano-2-methoxyiminoacetyl)-3-ethylurea

**5****USAGE and STORAGE FACTS:**

- Keep in its package closed.
- Exterminate the empty packages accordingly.
- Do not use the empty bottle for any other purpose.
- Do not refill the emptied food and beverage packages with the product.
- Toxic for the fish.
- Avoid contact with the underground or surface waters.

**STORING:**

Must be stored in a cool place, locked, and in its original package. If stored under these conditions, the product keeps its conformity to its specification for 2 years.

**COMPANY'S ANNOUNCEMENT:**

Agri Sciences guarantees the quality of this product provided that it is sold in its original package. All the damage that may result of the factors out of the company's control is at the buyer's risk. The company denies all the responsibility for the damages that may result of storing the product under wrong conditions or applying it in wrong amounts or at wrong times.

**AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.**

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79