

TIGANEX

Suspension Concentrate

Name and Amount of active ingredient: 300 g/L Pyrimethanil

N-(4,6-dimethyl pyrimidin-2-yl) aniline

Group: 9; D1, FUNGICIDE

READ THE LABEL BEFORE

DO NOT USE HOME

KEEP AWAY FROM CHILDREN, FOOD AND ANIMAL FOOD.

DO NOT BREATHE STEAM AND / OR PARTICLES

USE MASK, PROTECTIVE CLOTHING, GLOVES AND GLASSES

DO NOT EAT, DRINK OR SMOKE DURING THE APPLICATION

HUMAN AND ANIMAL ACCESS INTO THE AREA SPRAYED IS NOT PROHIBITED

TO PREVENT RISKS ON HUMAN AND ENVIRONMENTAL HEALTH FOLLOW INSTRUCTION MANUAL.

RECOMMENDED PRODUCTS OUT OF THE USE PROHIBITED.

Toxication Symptoms: Irritative for skin and mucosa. May cause headache, fatigue, diarrhea, cyanosis resulting of methemoglobina.

FIRST AID MEASURES:

Take the patient away from the sprayed area, take off the contaminated clothes.

Skin contact: Wash well with water and soap the areas of contact.

Eye contact: Rinse with running clean water for minimum 15 minutes keeping the eye lids open. Seek medical advice if the symptoms continue.

If inhaled: Take the patient out to clean air. Seek medical advice if the symptoms continue.

If swallowed: If the patient is conscious, make him vomit. Give him artificial respiration. Seek medical advice if the symptoms continue. Show him the product label and package.

Antidote: No specific antidote, symptomatic treatment is followed. Provide the patient oxygen. Use ascorbic acid or methylene blue.

PLANT	PEST	DOSE OF USE	TIME BETWEEN LAST SPRAYING AND HARVEST
Tomato	Gray mold, Botrytis cinerea	125 ml/100 L water	3 days
Grape	Gray mold, Botrytis cinerea	100 ml/100 L water	21 days
Apple	Venturia inaequalis, Apple scab	50 ml/100 L water	14 days

HOW TO USE:

Preparing:

Tiganex mixes easily with water. The necessary amount of product is mixed with water in a tank then put into the spraying tank the half of which is already filled with water while the mixer is on, and fill the rest with water. Pay attention for a good coating, especially the grape bunches must be well wetted

tomato: Spraying is started by the time the primal symptoms are observed, and repeated every 10-14 days according to the development of the disease. A good coating is essential.

Grape: Spraying is started by the time water begins to reach the grains. Repeated every 10-15 days according to the development of the disease. The rotation use with other fungicides with different effect prevents immunity.

Apple:

1st spraying; in red pimpernel period
2nd spraying; in pink bud stage
3rd spraying; when flower petals defoliate 70-80%
4th and further; with 10-14 days intervals according to climate conditions.

INFORMATION ON IMMUNITY:

Tiganex plant protection product has been classified in Group 9; D1 AS fungicide according to the mode of action. Repeated applications of plant protection products with the same mechanism of action, promotes the development of resistance. For this reason, to delay development of resistance, in the same growing season do not exceed the recommended total number of applications. Repeat the application if required, with a different mechanism of action (other than Group 9;D1).

Miscibility :

Cannot be used in mixture with aluminium containing products.

HAZARD FEATURE:

**N: DANGEROUS FOR THE ENVIRONMENT
CONTAINS PYRIMETHANIL**

RISK PHRASES:

R51: Very toxic to aquatic organisms.
R53: May cause long term negative effects in aquatic environments.

SAFETY PHRASES:

S1 / 2: Keep locked up and out of the reach of children.
S29: Do not empty into drains.
S56: This material and its package must be disposed at hazardous or special waste collection point.
S61: Avoid release into environment. Follow the special instructions of use and Safety Information sheet.

PRODUCER AND LICENSING COMPANY OWNER

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.
Bekir Saydam Cad. No:31 Pancar,Torbali-İZMİR
Tel: 0232 864 13 33 Fax: 0232 864 16 79

STORING CONSIDERATIONS :

- * Keep the plant protection product in a closed container.
- * Dispose empty packages properly.
- * Never use empty packages for any other purpose.
- * Do not refill empty food and beverage packages with plant protection products.
- * Toxic to fish.
- * Avoid infecting the underground and surface water.

STORAGE CONDITION:

Keep in its original package, tightly closed, away from the food, at a cool, and dry place, out of reach of children, under lock. From date of manufacture, if stored in original package, unopened, in a cool, dry condition, keeps its physical and chemical properties for at least 2 years.

COMPANY STATEMENT:

The manufacturer guarantees the quality of the product, provided that it is sold in manufacturer's original package. Damage which may occur beyond the control the manufacturer is at user's risk. Manufacturer is not responsible for any damage arising from improper storage conditions, or use of the product in wrong doses or at wrong times.

PRODUCER AND LICENSING COMPANY OWNER:

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31 Pancar,Torbalı-İZMİR

Tel: 0232 864 13 33 Fax: 0232 864 16 79