

Manufacture Date :
Expiration Date :
Charge No :
Permit Date & Number :

TARGETED PLANTS AND PESTS

PLANT NAME	APPLICATION METHOD & PERIOD	DOSAGE
VEGETABLES (Tomato, Pepper, Eggplant, Cucumber, Melon, Potato, Watermelon)	Seed Immersion: Immerse in pesticide solution for 8 hours before planting, or for 12 hours in case of potato tubers before cutting. For nurseries: Spray a few times once per 5 days. Before and after flowering: Spray plants before each flowering period.	17 ml/100 L water 9 ml/100 L water 17 ml/100 L water
FRUITS (Apple, Plum, Pear, Peach, Apricot, Citrus Fruits)	1. Sprout development: 2. Pink bud period 3. Fruits are pea sized.	17 ml/100 L water 17 ml/100 L water 17 ml/100 L water
VINEYARD	Application 1 and 2: New sprouts. Application 3: At the beginning of flowering period Application 4: Fine unripe period.	17 ml/100 L water 17 ml/100 L water 17 ml/100 L water
LEGUMES (Bean, Peas, Chickpea, Soy, Lentil, Peanut)	Spray a few times before and at the beginning of flowering stage in order to promote fruit formation.	20 -25 ml / da
CEREALS (Wetting seed on paddy, wheat, barley, corn)	Seed immersion: Immerse paddy seeds in pesticide solution for 12 hours before plantation. Pesticide should be sprayed at tillering, bolting and earing stages.	67 ml/100 L water 20 ml/da
STRAWBERRY	With 15 day intervals after flower emergence	75 ml/100 L water
COTTON	Application 1: At first ginning, Application 2: at first flowering Application 3: when 90% of bolls form	17 ml/100 L water 17 ml/100 L water 17 ml/100 L water
SUGAR BEET	Application 1: At the 2-3 leaved stage, Application 2: after thinning Application 3: after root thickening starts	9 ml/da 20 ml/da 20 ml/da
ORNAMENTAL PLANTS	Immerse bottom part of herbaceous plants for 5-10 min. Spray a few times with 10 day intervals. Spray a few times with one week intervals.	9 ml / 100 L water 6 ml / 100 L water 17 ml / 100 L water

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.



Name & Amount of Active Ingredient:

1- 3 g Sodium 5-nitroguaiacolate in Liter
2- 6 g Sodium ortho-nitrophenolate in Liter
3- 9 g Sodium para- nitrophenolate in Liter

GROUP	---	PGD
-------	-----	-----

FIRSTLY READ THE LABEL.
DO NOT USE AT HOME.
KEEP AWAY FROM CHILDREN, FOODSTUFFS AND ANIMAL FEED.
DO NOT INHALE VAPORS AND/OR PARTICLES OF PESTICIDE.
PLEASE USE PROTECTIVE CLOTHING, GLOVES AND GOGGLES.
DO NOT EAT, DRINK OR SMOKE DURING APPLICATION OF PESTICIDE.
DO NOT ALLOW HUMAN AND ANIMAL TO ENTRY INTO PESTICIDE TREATED AREAS FOR 1 DAY
PLEASE FOLLOW THE USAGE INSTRUCTIONS TO PREVENT ANY HUMAN HEALTH AND ENVIRONMENTAL RISKS.
IT IS STRICTLY PROHIBITED TO USE THE PESTICIDE ON ANYTHING OTHER THAN RECOMMENDED TARGETS.
SHAKE WELL AFTER ADDING CLEAN WATER TO EMPTY CONTAINER OF PESTICIDE WITH THE RATE OF 1/4. POUR THE SHAKEN WATER INTO DISINFECTION TANK. REPEAT THIS PROCESS THREE TIMES

NATIONAL TOXIN HOTLINE (UZEM) PHONE: 114

Net Amount :
Max. Retail Sales Price :

HAZARD TYPES:

Xi: Irritant

HAZARD SYMBOLS



Xi: Irritant

INGREDIENTS:

1- Sodium 5-nitroguaiacolate in Liter
2- Sodium ortho-nitrophenolate in Liter
3- Sodium para- nitrophenolate in Liter

RISK EXPRESSIONS RELATED TO APPLICATOR AND ENVIRONMENT:

R20/21/22: Harmful if inhaled, contact with skin or swallowed.

SAFETY EXPRESSIONS:

S2: Keep out of the reach of children
S13: Keep away from food, drink and animal feeding stuffs.
S29: Do not empty into drains
S36/37: Wear suitable clothes, protective clothes.
S46: Call a doctor immediately if swallowed and show the container or label.

MANUFACTURING AND PERMIT HOLDING COMPANY:

AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.
Bekir Saydam Cad. No: 31
Pancar, Torbalı, İzmir
Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

METHOD OF APPLICATION:

Effect Mechanism: It is easily absorbed into plant metabolism and increases plant sap mobility. Provides life force to all cells of the plant and positively affects and expedites germination, rooting, development and ripening processes. It especially has a positive effect on germination.

Properties: The product is a promoter that provides faster and more regular germination, stronger and faster development, and better and higher quality produce in all plants, applicable in all life stages of the plant from seed to harvest. It also acts as a stimulant protecting the plants against various adverse developmental conditions and activating biochemical and physiological resources of the plants.

Preparation:

Mix the recommended dosage of product with a small amount of water in a separate container. Fill the applicator tank with water to the half point and pour the pre-mixture in. Top off the tank with water thoroughly mix. Proceed to application in a timely fashion.

INFORMATION ON RESISTANCE:

The pesticide name **Sipgran** is a Plant Growth Designator classifies as Group -- in accord with its action mechanism. Repeated applications of pesticides with same action mechanisms invite development of resistance. Therefore, to delay development of resistance, do not exceed the recommended number of **Sipgran** applications in the same season. In case repeated application is necessary, take care to use pesticides with a different action mechanism (except of Group --).

MISCIBILITY:

It can be mixed with other pesticides and leaf fertilizer except of the ones having very-strong acid characters

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

**Name & Amount of Active Ingredient:**

1- 3 g Sodium 5-nitroguaiacolate in Liter
2- 6 g Sodium ortho-nitrophenolate in Liter
3- 9 g Sodium para- nitrophenolate in Liter

GROUP	---	PGD
--------------	------------	------------

SYMPTOMS OF POISONING:

Headache, sweating, vomiting, weakness, nausea.

FIRST AID MEASURES:

May be harmful if swallowed. Therefore, drink plenty of water and milk.

In case of contact with skin, wash the contaminated parts with plenty of water and soap.

ANTIDOTE AND NECESSARY INFORMATION:

There is no special antidote. The treatment is performed as per the symptoms.

NATIONAL TOXIN HOTLINE (UZEM) PHONE: 114

HANDLING AND STORAGE CONSIDERATIONS:

- Store the pesticide in its closed packaging.
- Suitably dispose the empty packages.
- Do not use empty packages for any other purpose.
- Do not put pesticide in empty food or drink containers.

STORAGE CONDITIONS:

In its original package store it closed , away from food and Keep locked up and out of the reach of children and in cool and dry place. When stored under cool and dry conditions in its original closed package, the product will not experience any physical, chemical or chemical changes outside the tolerance limits for 2 years.

DISCLAIMER:

The manufacturer warrants the quality of the product as long as it is sold in its original packaging. The user is responsible for the damage risk, arising from the factors out of the manufacturer. The manufacturer shall not accept any responsibility regarding damages originating from erroneous storage or failure to follow instructions.

MANUFACTURING AND PERMIT HOLDING COMPANY:

AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.
Bekir Saydam Cad. No: 31
Pancar, Torbali, Izmir
Tel: +90-232-864-13-33 Fax: +90-232-864-16-79