

Manufacture Date :
Expiration Date :
Batch No :
Registration Date & Number :

TARGETED PLANTS AND PESTS

PLANT NAME	PEST NAME	DOSAGE & STAGE	LAST APPLICATION-HARVEST INTERVAL
COTTON	Two-dotted red spider (<i>Tetranychus urticae</i>)	40 ml/da egg, nymph, mature	21 days
	Cotton red spider (<i>Tetranychus cinnabarinus</i>)	40 ml/da egg, nymph, mature	
AUBERGINE	Two-dotted red spider (<i>Tetranychus urticae</i>)	50 ml/da nymph, mature	3 days
	Cotton red spider (<i>Tetranychus cinnabarinus</i>)	50 ml/da nymph, mature	
WATERMELON	Two-dotted red spider (<i>Tetranychus urticae</i>)	50 ml/da nymph, mature	3 days
	Cotton red spider (<i>Tetranychus cinnabarinus</i>)	50 ml/da nymph, mature	
TOMATO	Two-dotted red spider (<i>Tetranychus urticae</i>)	50 ml/100 L water, nymph, mature	3 days
	Cotton red spider (<i>Tetranychus cinnabarinus</i>)	50 ml/100 L water, nymph, mature	
	White fly (<i>Bemisia tabaci</i>)	60 ml /100 L water, larva, nymph	
STRAWBERRY	İki noktalı kırmızı örümcek (<i>Tetranychus urticae</i>)	50 ml/100 L water, nymph, mature	3 days
	Cotton red spider (<i>Tetranychus cinnabarinus</i>)	50 ml/ water, nymph, mature	

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



Suspension Concentrate (SC)

Name & Amount of Active Ingredient: in Liter 240 g
Spiromesifen 2-oxo-3-(2,4,6-trimethylphenyl)-1-oxaspiro[4.4]non-3-en-4-yl 3,3-dimethylbutanoate

GROUP	23	ACARICIDE
-------	----	-----------

PLEASE READ THE LABEL FIRST.
DO NOT USE AT HOME.
KEEP AWAY FROM CHILDREN, FOODSTUFFS AND ANIMAL FEED.
DO NOT INHALE VAPORS OF PESTICIDE.
PLEASE USE MASK, PROTECTIVE CLOTHING, GLOVES AND GLASSES.
DO NOT EAT, DRINK OR SMOKE DURING APPLICATION OF PESTICIDE.
DO NOT PERMIT HUMAN AND ANIMAL ENTRY INTO PESTICIDE TREATED AREAS FOR 12 HOURS.
PLEASE FOLLOW THE USAGE INSTRUCTIONS TO PREVENT ANY HUMAN HEALTH AND ENVIRONMENTAL RISKS.
IT IS STRICTLY PROHIBITED TO USE THE PESTICIDE ON ANYTHING OTHER THAN RECOMMENDED TARGETS.

NATIONAL TOXIN HOTLINE (UZEM) PHONE:114

Net Amount :
Max. Retail Sales Price :

HAZARD TYPES:

Xi: Irritant N: Dangerous for environment

HAZARD SYMBOLS



Xi: Irritant



N: Dangerous for environment

INGREDIENTS:

CONTAINS Spiromesifen.

RISK EXPRESSIONS AND DETAILS:

R43: It may constitute sensitivity in case of contact with skin.
R50: Very toxic to aquatic organisms
R53: May cause long-term adverse effects in the aquatic environment

GÜVENLİK İBARESİ VE AÇIKLAMASI:

S1/2: Keep locked up and out of the reach of children
S13: Keep away from food, drink and animal feeding stuffs
S29: Do not empty into drains
S36/37/39: Wear suitable protective clothing, gloves and eye/face protection
S46: If swallowed, seek medical advice immediately and show this container or label
S56: Dispose of this material and its container at hazardous or special waste collection point
S61: Avoid from releasing it to environment. Apply Special Use Instruction/Safety Data Sheet.

MANUFACTURING AND PERMIT HOLDING COMPANY:

AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.

Bekir Saydam Cad. No: 31
Pancar, Torbali, Izmir, TURKEY
Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

METHOD OF USE:**Preparation of Disinfection:**

Firstly, required pesticide is mixed with some water in a container. The prepared mixture is added to the disinfection device tank that is full in half level with water. The tank is filled with water again and mixed. It is ready to perform disinfection. This mixture is sprayed all parts of the plant when the weather is calm and cool.

Disinfection Time:

Cotton: Separate the field in 4 pieces. In each part, take one leaf from 25 plants in every 10-15 step. Count on 100 leaves via a magnifying glass totally. The disinfection is performed if there are Red Spiders with the number of 5 in Mediterranean, 10 in Aegean and Southeast per leaf. Partial disinfection may be performed in case of side contamination. The water quantity for a decare must be determined via calibration according to the device to be used. And then required pesticide must be added. If tractor-mounted- ground devices will be used, inter-row nozzles must be used.

Tomato, Aubergine, Watermelon: Beginning from the first inflorescence, enter the field crossly from the edge. Count with magnifying glass totally 10-20 leaves by one leaf from bottom and intermediate leaves once in 3-5 steps. The disinfection is performed if there are 3 or 5 Red Spiders per leaf. The water quantity for a decare must be determined via calibration according to the device to be used. And then required pesticide must be added.

Strawberry: The disinfection is performed when 15 moveable things per leaf are visible from 50 leaves representing the field. If there are ripe fruits, obey the waiting time or perform the disinfection after collecting the fruits.

White fly: The disinfection is performed when you see 0.5-2 larva in May-June period; 2-3 larva in July-August period and 3 larva in September per leaf considering 50 leaves.

INFORMATION ON RESISTANCE:

The pesticide name **Koffex** is a fungicide classifies as Group 23 in accord with its action mechanism. Repeated applications of pesticides with same action mechanisms invite development of resistance. Therefore, to delay development of resistance, do not exceed the recommended number of **Koffex** applications in the same season. In case repeated application is necessary, take care to use pesticides with a different action mechanism (other than Group 23).

MISCIBILITY:

It may be mixed with many insecticides, fungicide and leaf fertilizer. But, pre-mixture test must be performed before the application.

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.

**Suspension Concentrate (SC)**

Name & Amount of Active Ingredient: 1n Liter 240 g
Spiromesifen 2-oxo-3-(2,4,6-trimethylphenyl)-1-oxaspiro[4.4]non-3-en-4-yl 3,3-dimethylbutanoate

GROUP	23	ACARICIDE
--------------	-----------	------------------

PUT CLEAN WATER TO THE EMPTY CONTAINER OF THE PESTICIDE TO THE LEVEL OF ONE QUARTER AND SHAKE IT. POUR THE SHAKEN WATER INTO THE DISINFECTION TANK. PERFORM THIS APPLICATION FOR 3 TIMES.

“THIS PESTICIDE ARE NOT USED ON TOMATO”

SYMPTOMS OF POISONING:

There are no specific symptoms of poisoning on human.

FIRST AID MEASURES:

In case of contact with the skin, wash it with plenty of water. Take off the contaminated clothes. In case of contact with eye, wash with plenty of water for at least 15 minutes. Call a doctor. Take the label and package with you.

ANTIDOTE AND IMPORTANT INFORMATION:

There is no special antidote. In accord with symptoms supportive treatment applied.

NATIONAL TOXIN HOTLINE (UZEM) PHONE:114

HANDLING AND STORAGE CONSIDERATIONS:

- Store the pesticide in its closed packaging.
- Suitably dispose the empty packages.
- Do not use empty packages for any other purpose.
- Do not put pesticide in empty food or drink containers
- Not toxic for mature bees and bumble bee.
- Toxic for fish.
- Avoid contaminating above- and underground waters.

STORAGE CONDITIONS:

In its original package store it closed , away from food and Keep locked up and out of the reach of children and in cool and dry place.

When stored under normal (cool and dry) conditions in its original closed package, the product will not experience any physical, chemical or chemical changes outside the tolerance limits for 2 years.

DISCLAIMER:

The manufacturer warrants the quality of the product as long as it is sold in its original packaging. The manufacturer shall not accept any responsibility regarding damages originating from erroneous storage or failure to follow instructions.

WARNING: Shake well before use.

MANUFACTURING AND PERMIT HOLDING COMPANY:

AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.

Bekir Saydam Cad. No: 31

Pancar, Torbali, Izmir, TURKEY

Tel: +90-232-864-13-33 Fax: +90-232-864-16-79