

1

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

NICARDEN

Powder Wettable in Water

Active Ingredient Name and Ratio:70% Propineb

Polymeric zinc 1,2-propylenebis (dithiocarbamate)

GROUP:M3, FUNGICIDE

CAUTION

READ THE LABEL FIRST.

DO NOT USE AT HOME.

KEEP AWAY FROM CHILDREN, FOOD AND ANIMAL FOOD.

DO NOT INHALE THE VAPOR AND/OR DRIBLETS.

WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES.

DO NOT EAT, DRINK, OR SMOKE DURING THE APPLICATION.

AREA APPLIED IS NON-ACCESSIBLE FOR HUMANS AND ANIMALS FOR 14 DAYS.

FOLLOW THE USING INSTRUCTIONS IN ORDER TO PREVENT THE RISKS AGAINST HUMAN AND ENVIRONMENT HEALTH.

THE USE ON THE PRODUCTS OTHER THAN RECOMMENDED STRICTLY PROHIBITED.

TOXICATION SYMPTOMS:

Despite its low toxicity, if used uncarefully or swallowed bu mistake, may result in dizziness, nausea, and vomitting.

FIRST AID PRECAUTIONS:

Call the doctor, let him see the product label and package. Take the patient away from the sprayed area. Take the contaminated uniform off, wash well with soap and water if contacted skin, and eye. If swallowed, the patient is made vomit.

ANTIDOTE: No any special antidote.

NATIONAL TOXIC CONSULTANCY ASSOCIATION (NPCA) Tel:114

2

Production date:

Expiry date:

Charge nr:

License date and nr:

PLANTS and DISEASES TO APPLY ON

| Plant | Disease | Dosage and Timing | Period between final spraying and harvest | | |
|----------|-------------------------------------|-------------------|---|--|--|
| rutaceae | alternaria alternata f. spp. citri) | 250 g/100 l water | 7 days | | |
| apple | venturia inaequalis | 200 g/100 l | 7 days | | |
| pear | venturia pirina | 250 g/100 l | 7 days | | |
| orchard | plasmopara viticola | 200 g/100 l | 28 days | | |

| | | | | | |
|---------------------------------|---|--|----------------------|--|--|
| | | | | | |
| pistacchio | septoria pistacina | 300 g/100 l water | 28 days | | |
| *vegetable seeds | fusarium spp, pythium spp, rhizoctonia spp, | 200-250 g/100 l water | 7 days | | |
| tomato | cladosporium fulvum | 300 g/100 l water | 7 days | | |
| | phytophthora infestans | 200 g/100 l water | 7 days | | |
| tomato, potato, aubergine | alternaria solani | 300 g/100 l water | 7 days | | |
| potato | phytophthora infestans | 200 g/100 l water | 7 days | | |
| beans | colletotrichum spp Uromyces phaseoli | 200 g/100 l water 200 g/100 l water | 7 days 7 days | | |
| chickpeas | ascochyta rabiei | 200 g/100 l water | 7 days | | |
| cucumber | pseudoperonospora cubensis | 200 g/100 l water | 7 days | | |
| onion | peronospora destructor | 200 g/100 l water | 7 days | | |
| cabbage | peronospora brassicae | 200 g/100 l water | 7 days | | |
| lettuce | bremia lactucae | 200 g/100 l water | 7 days | | |
| melon- watermelon | colletotrichum spp. | 200 g/100 l water | 7 days | | |

| | | | | | |
|-----------|--------------------------|-------------------|---------|--|--|
| | | | | | |
| hop | pseudoperonospora humuli | 150 g/100 l water | 35 days | | |
| safflower | alternaria spp. | 200 g/100 l water | 35 days | | |
| carnation | Uromyces caryophyllus | 200 g/100 l water | ---- | | |
| rose | Phragmidium spp. | 200 g/100 l water | ---- | | |

(*)The dose varies according to the level of the disease and the plant vegetation duration.

3

HOW TO USE

Preparing for applying:

The tank of the spraying tool is filled with water in half. The necessary amount of the product is mixed with some amount of water in another tank and put into the tank. Water is added until the tank is completely full while the mixer is on. Must be made at the early times or cool evening times of the day when it is not windy. In case of spraying, coating must be done well, the back sides and also the whole parts as well of the plant must be sprayed very well.

Using of the Product

venturia inaequalis:1. spraying, by the time the flower buds pop, 2. spraying, at the pink-white badge bud term, 3. spraying by the time 70-80% of the petals fall, 4. and the others, according to the weather conditions and the development of the disease.

plasmopara viticola:1 spraying by the time the buds are 25-30 cm long, 2. and the others according to the development of the disease and the meteorologic conditions, if the convenient conditions for infection are present, with intervals of 8-10 days. The spraying is stopped by the time such conditions do not exist anymore.

phytophthora infestans:By the time the primal marks on the tomato leaves are noticed, the spraying is started. Repeated each 7-10 days according to the degree of the disease and the weather conditions.

colletotrichum spp:Spraying is started by the time the primal symptoms are noticed around. Repeated each week according to the development of the disease and the weather conditions.

INFORMATION ON IMMUNITY:

The crop protect product Nicarden is a fungicide that is grouped as M3 according to its effect mechanism. The repetitive applications of the plant protect products with the same effect

mechanism encourages the development of immunity. For this reason, in order to delay the development in immunity, do not exceed the advised total number of application within the same production period. In case the application should be repeated, try to use of the plant protect products with different mechanism effect (except the Group M3).

MISCIBILITY

Can be mixed with Cooperoxychlorure.

4

Net Amount:

Maximum Retail Sales Price:

DANGER PROPERTY

T:TOXIC N:DANGEROUS FOR THE ENVIRONMENT

DANGER SYMBOLS

T:TOXIC N:DANGEROUS FOR THE ENVIRONMENT

INGREDIENTS:CONTAINS PROPINEB

JARGONS TO RISK AND EXPLANATIONS:

R20:Hazardous to inhale

R43:May cause skin irritation.

R 48/20722:Hazardous:Serious damage on health if inhaled for long and swallowed.

R50:Very toxic for the aquatic organisms.

JARGON TO SAFETY AND EXPLANATIONS

S2:Keep out of the reach of children.

S 36/37:Use appropriate protective uniform and gloves.

S22:Do not inhale the powders.

S24:Avoid contact with skin.

S46:If swallowed, immediately see a doctor, let him see the package or the label.

S61:Avoid releasing it into the environment. Address to the Special Using Instructions/Security Information Form.

CAUTIONS ON USING AND STORING:

*Keep closed it its package.

*Exterminate the emptied packages accordingly.

*Do not use the emptied packages for any other purpose.

*Do not refill the emptied food and beverage packages with the product.

*Toxic for the fish.

*Do not contaminate into the under ground and surface waters.

5

STORING:

Keep in the original package, closed, away from food and out of the reach of children, and animals, at dry and cool places, under lock. Keeps its physical and chemical properties min. for 2 years after the production date if stored under cool and dry conditions, in the original, not opened package.

COMPANY'S ANNOUNCEMENT:

The company guarantees quality provided that the product is sold in original package.

All the damage that may occur of the factors out of the control of the manufacturer are at the user's risk. The company denies all the responsibility for the damages that may result of storing the product under wrong conditions or applying it at wrong doses and times.

MANUFACTURER AND LICENSED COMPANY:

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79