

Manufacture Date :
 Expiration Date :
 Charge No :
 Permit Date & Number :

TARGETED PLANTS AND PESTS

PLANT NAME	PEST NAME	DOSAGE	LAST APPLICATION-HARVEST INTERVAL
CITRUS	Against red scale (<i>Aonidiella citrina</i>)	50 ml/ 100 L water larva	28 days
CUCUMBER	White fly (<i>Bemisia tabaci</i>)	50 ml/da/ larva, pupa	3 days
COTTON	White fly (<i>Bemisia tabaci</i>)	50 ml/da/ larva, pupa	28 days
PEPPER (greenhouse)	White fly (<i>Bemisia tabaci</i>)	50 ml/ 100 L water Larva, pupa	3 days
TOMATO	White fly (<i>Bemisia tabaci</i>)	50 ml/ 100 L water Larva, pupa	1 day
PEACH	Mulberry scale (<i>Pseudaulacaspis Pentagona</i>)	50 ml/100L water	14 days

PROPERTIES:

Product holds its pest control effect for 20 to 30 days. Eggs already on the plant before application, as well as eggs laid after application, cannot hatch. Product prevents development of larvae and nymphs, and kills them during their life stage shift.

METHOD OF USE:

Preparation:

First mix the prescribed amount of pesticide with a small amount of water. Then fill half of the reservoir of the sprayer with water. Then pour the mixture into reservoir. Keep mixing the reservoir content while you top up the reservoir with water.

Application:

Application should be done after morning dew evaporates or during evening cool. The product can be applied by any type of sprayer or atomiser.

Against Red Scale: Application should be done when winter adults start to breed and most of the pests are in the first stage.

White fly: Application should be done when 5 white fly larvae per leaf are observed.

White mulberry scale: Application against first and second spawns should be done at the time of first larvae hatching, and again after a 20 day interval.

Continued on Page 2

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



Emulsifiable Concentrat (EC)

Name & Amount of Active Ingredient: 100 g/L Priproxyfen 4-phenoxyphenyl (RS)-2-(2-pyridyloxy) propylether

GROUP	7C	FUNGICIDE
-------	----	-----------

FIRSTLY READ THE LABEL.
 DO NOT USE AT HOME.
 KEEP AWAY FROM CHILDREN, FOODSTUFFS AND ANIMAL FEED.
 DO NOT INHALE VAPORS AND/OR PARTICLES OF PESTICIDE.
 PLEASE USE PROTECTIVE CLOTHING, GLOVES AND GLASSES.
 DO NOT EAT, DRINK OR SMOKE DURING APPLICATION OF PESTICIDE.
 NO LIMITATION REQUIRED FOR HUMAN AND ANIMAL ENTRY INTO PESTICIDE TREATED AREAS FOR 4 HOURS.
 PLEASE FOLLOW THE USAGE INSTRUCTIONS TO PREVENT ANY HUMAN HEALTH AND ENVIRONMENTAL RISKS.
 IT IS STRICTLY PROHIBITED TO USE THE PESTICIDE ON ANYTHING OTHER THAN RECOMMENDED TARGETS.

Symptoms of Poisoning: nausea, vomit, diarrhoea.
First Aid Measures: If contact with the medicine, take off contaminated clothes. Wash the skin part with plenty of water. Wash with plenty of water in case of contact with the eye. If drug is swallowed, call the doctor immediately. If swallowed, drank sip by sip 1 glass to the patient. If the doctor doesn't say, do not vomit the patient. Be not given anything to unconscious patient.
Antidote and Necessary Information: There is no special antidote. Treatment is applied according to the symptoms.

NATIONAL TOXIN HOTLINE (UZEM) PHONE: 114

Net Amount :
 Max. Retail Sales Price :

HAZARD TYPES:

Xn: Harmful
 N: DANGEROUS FOR THE ENVIRONMENT

HAZARD SYMBOLS:



Xn: Harmful



N: DANGEROUS FOR THE ENVIRONMENT

INGREDIENTS:

CONTAINS PYRIPROXYFEN AND KSILEN

RISK EXPRESSIONS:

R10: it inflames
R20/21: It is harmful in case of inhaled and contact with skin.
R38: It irritates to skin.
R50: It is too toxic for aquatic environments.
R53: may cause long-term negative effects in aquatic environments.
Safety Phrases:
S1/2: Lock it up and keep out of the reach of children.
S13: Keep away food, drinks and animal feeds.
S29: Do not empty into drains.
S43: If fired, use pressured water, CO2, fire extinguisher powder, foam, sand.
S36/37/39 : Wear suitable protective clothing, gloves and eye/face protection.
S43: If fired, use pressured water, CO2, fire extinguisher powder, foam, sand.
S56: Dispose this material and the container in dangerous and special waste collecting places.
S64: In case of swallow, wash the mouth with water (If the patient is conscious)

MANUFACTURING AND PERMIT HOLDING COMPANY:
 AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.

Bekir Saydam Cad. No: 31
 Pancar, Torbali, Izmir, TURKEY
 Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

(Company Stamp & Signature)

INFORMATION ON RESISTANCE:

The pesticide name **Pyrabiol** is an acaricide classified as GroupUN in accord with its action mechanism. Repeated applications of pesticides with same action mechanisms invite development of resistance. Therefore, to delay development of resistance, do not exceed the recommended number of **Pyrabiol** applications in the same season. In case repeated application is necessary, take care to use pesticides with a different action mechanism (except form Group UN).

MISCIBILITY:

Except for alkaline characteristics, it can be mixture with common fungicide, insecticide and foliar fertilizers.

AGRI SCIENCES TARIM ve ILAC UR. SAN. ve TIC. LTD. STI.



INTRACURE

Emulsifiable Concentrât (EC)

Name & Amount of Active Ingredient: 100 g/L Priproxyfen
4-phenoxyphenyl (RS)-2-(2-pyridyloxy) propylether

GROUP	7C	FUNGICIDE
--------------	-----------	------------------

HANDLING AND STORAGE CONSIDERATIONS:

- Store the pesticide in its closed packaging.
- Suitably dispose the empty packages.
- Do not use empty packages for any other purpose.
- Do not put pesticide in empty food or drink containers.
- Toxic for fish
- Avoid contaminating above- and underground waters.

STORAGE CONDITIONS:

In its original package store it closed , away from food and keep locked up and out of the reach of children and in cool and dry place. When stored under normal (cool and dry) conditions in its original closed package, the product will not experience any physical, chemical or chemical changes outside the tolerance limits for 2 years.

DISCLAIMER:

The manufacturer warrants the quality of the product as long as it is sold in its original packaging. User endures damage risk which can occur because of factors out of manufacturer control. The manufacturer shall not accept any responsibility regarding damages originating from erroneous storage or failure to follow instructions.

MANUFACTURING AND PERMIT HOLDING COMPANY:

AGRI SCIENCES TARIM VE ILAC UR. SAN. VE TIC. LTD. STI.
Bekir Saydam Cad. No: 31
Pancar, Torbali, Izmir, TURKEY
Tel: +90-232-864-13-33 Fax: +90-232-864-16-79

(Company Stamp & Signature)