

Production Date :  
Expire Date :  
Batch Number :  
License Date and Number. :

**PLANTS AND PESTS TO BE USED FOR**

| Plant                 | PEST  | USAGE DOSAGES    | WAITING PERIOD |
|-----------------------|---|------------------|----------------|
| Cucurbitaceae (Melon) | Powdery Mildew (Erysiphe cichoracearum, Sphaerotheca fuliginea) | 15 g/100 L water | 3 days         |
| Vineyard              | Powdery Mildew (Uncinula necator)                               | 10g/ 100 l water | 14 days        |
| Apple                 | Codling Moth (Venturia inaequalis)                              | 15 g/100 L water | 14 days        |
| Circuit               | alternaria solani (Alternaria alternata f.sp.citri)             | 20 g/100 L water | 28 days        |
| Rice (Paddy)          | Paddy Rot (Pyricularia oryzae)                                  | 20g/da           | 35 days        |

**USAGE:**

**Preparing the pesticide for use:**

First, adequate amount of pesticide is poured into water in a different cup and waited 1-2 minutes for precipitation. After that it is mixed until the pesticide dissolves completely. Prepared mixture is poured in to the reservoir of the disinfection tool which is half-filled with water. The reservoir is filled with adequate amount of water.

**Melon:** Disinfection should be started before the disease observed and in the occasion that proper conditions exist for disease, disinfection should be repeated with 7-10 days intervals.

**Vineyard:**

1. Disinfection when offshoots are 25-30 cm long
2. Disinfection when grapes are as big as pellets
3. Disinfection after 15 days after second disinfection and subsequent disinfections are continued once per 15 day.

**Apple Blackspot Bruising:**

1. Disinfection when the flower cells are swelled
2. Disinfection when pink flower colliculus period
3. Disinfection when 70 – 80 % of the flower petals are fallen. Subsequent disinfections are repeated with 10-12 day intervals if the conditions are still suitable for the disease.

**AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.**



**RAPIDOX 50 WG**

Water Dispersible Granule (WP)

**Name and quantity of the active component:** %50 Trifloxystrobin methyl (E)-methoxyimino-(E)-a(1-(a,a,a-trifluoro-m-tolyl)ethylideneminoxy)-o-tolyl)acetate

|              |           |                  |
|--------------|-----------|------------------|
| <b>GROUP</b> | <b>11</b> | <b>FUNGICIDE</b> |
|--------------|-----------|------------------|

READ THE LABEL FIRST.

DO NOT USE AT HOME.

KEEP OUT OF REACH OF CHILDREN, FOOD AND ANIMAL FEED.

AVOID INHALATION OF VAPORS OR PARTICULES.

WEAR MASK, PROTECTIVE CLOTH, GLOVES, GOGGLES.

DO NOT EAT, DRINK OR SMOKE DURING TREATMENT.

DO NOT USE THE MEDICATED SEEDS AS FOOD OR ANIMAL

FEED.

OBEBY INSTRUCTIONS FOR USE TO AVOID THE RISK OF HUMAN AND ENVIRONMENTAL HEALTH.

USING TOGETHER WITH THE PRODUCTS NOT ADVISED CERTAINLY FORBIDDEN.

**TOXICOLOGICAL SYMPHOMS:** There is no symptom known.

**FIRST AID MEASURES:** If you suspect for poisoning, break out immediately, call a doctor, show label. If swallow and the patient is conscious, vomiting him. Give him a plenty of active carbon and treatment symptomaticly. If the patient isn't conscious, don't give anything via oral.

**ANTIDOTE AND REQUIRED INFORMATION:** No specific antidote is known. Treatment symptomaticly.

**NATIONAL POISON INFORMATION OFFICE: 114**

Net Weight :  
Maximum Selling Price:

**HAZARD PROPERTY**

Xi: Irritation N: Harmful for the Environment

**DANGER SYMBOLS**



Xi: Irritation N: Harmful for the Environment

**INGREDIENTS:**

**It contains trifloxystrobin.**

**RISK PHRASES AND DISCLOSURES**

**R43** May cause sensitization by skin contact.

**R50/53** Very toxic to aquatic organisms. May cause long-term adverse effects in aquatic environments.

**SAFETY PHRASES AND DISCLOSURES**

**S1/2** Keep out of reach of children

**S13** Keep away from foods, drinks and animal feedstuffs

**S24** Avoid to contact skin.

**S36/37/39** Use suitable protective clothing, gloves and protective goggles/mask.

**S46** If swallowed, seek medical advice immediately and show the label.

**S56** Dispose the material at hazardous or special waste collection point.

**PRODUCER AND OWNER OF THE LICENCE:**

**AGRI SCIENCES Tarım ve İlaç. Ür.**

San. ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31 Pancar Torbalı, İZMİR

Tel: 0090 0232 864 13 33

Fax: 0090 0232 864 16 79

**Citrus:** Disinfection should be started in the spring when the offshoots become 5-10 cm long and the first indications of disease is observed in the new leaf. If necessary disinfection should be repeated with 12-14 day intervals.

**Rice:** First disinfection should be applied when the first disease indications are observed. If the conditions are convenient for disease, other disinfections should be applied once per 10-15 day.

**INFORMATION ABOUT RESISTANCE:**

**Rapidox 50 WG** named plant protection products according to its mechanism of action is classified as Group 11. Repetitive applications of plant protection products with the same mechanism of action, encourages the development of resistance. For this reason, to delay development of resistance in **Rapidox 50 WG** on the same production season, the number of applications exceeds the recommended total. In cases where the application should be repeated, with a different mechanism of action (Group 11 external) careful use of plant protection products.

**MIXING CASES:** Generally, it can be mixed with most of the commonly used fungicides. In any hesitation, a mixing test should be done before using to full extent. If Rapidox 50 WG will be used with another pesticide in the same tank, Rapidox 50 WG should be poured into the tank first and then other pesticides should be pesticides after mixing with water in a different cup. If the other pesticides have a Water-Soluble ( WSB ) label, those pesticide should be poured to the reservoir before Rapidox 50 WG.

**Citrus:** Disinfection should be started in the spring when the offshoots become 5-10 cm long and the first indications of disease is observed in the new leaf. If necessary disinfection should be repeated with 12-14 day intervals.

**Rice:** First disinfection should be applied when the first disease indications are observed. If the conditions are convenient for disease, other disinfections should be applied once per 10-15 day.

**AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.**



# RAPIDOX 50 WG

Water Dispersible Granule (WP)

**Name and quantity of the active component:** %50 Trifloxystrobin methyl (E)-methoxyimino-(E)-a(1-(a,a,a-trifluoro-m-tolyl)ethylideneminoxy)-o-tolyl)acetate

| GROUP | 11 | FUNGICIDE |
|-------|----|-----------|
|-------|----|-----------|

**NATIONAL POISON INFORMATION OFFICE: 114**

**PLANT TOLERANCE**

Any phytotoxic occurs for recommended plants If use properly.

**SAFETY PRECAUTIONS DURING USING AND STORAGE:**

- \* Keep in closed original container
- \* Dispose of empty containers properly.
- \* Do not use empty containers for other purposes.
- \* Do not put food and drink boxes, empty plant protection product.
- \* Harmful for fishes
- \* Avoid infecting ground and surface water.

**STORAGE CONDITIONS:**

Store in dry and cool conditions, in closed original container separately from feed and food, If stored in cool and dry conditions, the product will keep its physical, chemical and biological structure at least 2 years.

**COMPANY DECLARATION:**

The quality of the pesticide is guaranteed by the producer if it is to be sold in the original packaging. The producer will not be responsible of the losses and damages resulting from the handling and storage mistakes or wrong treatment or improper actions not mentioned on the label.

**PRODUCER AND OWNER OF THE LICENCE:**

**AGRI SCIENCES Tarım ve İlaç. Ür.**  
San. ve Tic. Ltd. Şti.  
Bekir Saydam Cad. No:31 Pancar Torbalı, İZMİR  
Tel: 0090 0232 864 13 33  
Fax: 0090 0232 864 16 79