

1

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

# AGRISAVE

## Solution Concentrate (SL)

Active Ingredient:722 g Propamocarb-HCL /L

Propyl 3-(dimethylamino) propylcabamate hydrochloride

**GROUP: 28; F4; FUNGICIDE**

### CAUTION

READ THE LABEL FIRST.

DO NOT USE AT HOME.

KEEP AWAY FROM CHILDREN FOOD, AND ANIMAL FOOD.

DO NOT INHALE THE VAPOR AND DRIBLETS

WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES

DO NOT EAT, DRINK, OR SMOKE DURING THE APPLICATION

AREA APPLIED IS NON-ACCESSIBLE FOR HUMANS AND ANIMALS FOR A DAY

FOLLOW THE USING INSTRUCTIONS IN ORDER TO AVOID RISKS ON HUMAN AND ENVIRONMENT HEALTH

THE USE ON THE PRODUCTS OTHER THAN RECOMMENDED STRICTLY PROHIBITED

### TOXICATION SYMPTOMS:

Headache, dizziness, fatigue, temper loss, fever.

### FIRST AID PRECAUTIONS:

Consult a doctor and show him the product package or label. Keep the patient away from the area of effect, take off the contaminated uniforms immediately, wash well the skin surfaces that contacted with water and soap. If swallowed, give patient salty water to make him vomit.

### ANTIDOTE AND TREATMENT:

If swallowed so much that it prevents cholinesterase, ATROPINE SULPHATE , is the antidote. If swallowed in small quantity, symptomatic treatment is followed.

NATIONAL TOXIC CONSULTANCY ASSOCIATION (NPCA) Tel:114

## 2

Production date:

Expiry date:

Charge nr:

License date and nr:

PLANTS and HAZARDOUS ORGANISMSTO APPLY ON

<b>Plant</b>	<b>Disease</b>	<b>Dosage and Timing</b>	<b>Period between final spraying and harvest</b>		
sunflower	plasmophora helianthi	500 ml/100 kg seed	-		
cucurbitacea	Pseudoperonospora cubensis	250 ml/da	3 days		
potato	Phytophthora infestans	250 ml/da	3 days		
tobacco	Phythium spp.	7,5 ml/m2	3 days		

**HOW TO USE:**

First, make a mixture with water in a separate tank. Fill in the half of the tool tank, and add the mixture into it, while the mixer is on. Then add more water in the necessary amount while the mixer is on.

**On seeds:** Shall directly be smudged onto the seeds.

**Sunflower, plasmophora helianthi:** Spraying is made before planting as seed spraying.

**Green nozzle spraying;**

**Pseudoperonospora cubensis, cucurbitacea:** Spraying is started by the time the plants start branching or the primal symptoms are noticed around, and repeated every 7-10 days according to the grade of the disease and the climate conditions. There must be one week between the last spraying and harvest.

**Phytophthora infestans, potato:** Spraying is started on the areas where the disease is seen each year, before the disease starts by the time the daily average temperature reaches 16°C and the lowest temperature is 10°C. On the areas where the disease is not seen each year, the spraying is started by the time the primal disease symptoms are noticed around or within the field. Within the regions where the weather is dry and hot, 2-3 applications with intervals of 14 days, within the regions where it is cool and rainy repetitive applications according to the development of the disease are made.

**Phythium spp., tobacco:** Spraying is done after the seeds are planted, and cover fertilizer is spreaded onto, or after the emergence of the seedlings is over and the disease is seen.

### 3

**IMMUNITY:**

Agrisave is a fungicide classified as Group 28, F4 according to its effect mechanism. Repetitive applications of the plant protect products with the same effect mechanism encourages a development of immunity. In order to delay this development, do not exceed the total number of application amount of Agrisave advised for the season. In case a repeat is needed, use plant protect product with different effect mechanism (out of the Group 28,F4).

**MISCIBILITY:**

Can be used with other fungicides as a tank mixture. Cannot be mixed with bordeaux mixture and foliar fertilizers.

When used for seeds, it must be used solely without being mixed with anything else.

### 4

**Net Amount:**

**Min. Retail Sales Price:**

DANGER:Xi Sensitizer

Contains Propamocarb-HCL

## **RISK JARGON AND EXPLANATIONS**

R43:May irritate skin in case of contact.

## **SAFETY JARGON AND EXPLANATIONS**

S24:Avoid skin contact

S36/37:Wear appropriate protective uniform, and gloves

## **USAGE and STORAGE FACTS:**

- Keep the product in its package, closed.
- Exterminate the emptied bottles accordingly.
- Do not use the emptied packages for any other purposes.
- Do not refill the emptied food and beverage cans with the product.
- Non-toxic for the bees and fish. Avoid contaminating into the underground and surface waters.

## **MANUFACTURER AND LICENSED COMPANY:**

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79

# **5**

## **STORING:**

Keep in its original package, closed, away from food, out of the reach of children, under lock. If stored under normal conditions (cool and dry), in original package, and closed, keeps its physical, and chemical properties for min. 2 years after the production date.

## **COMPANY'S ANNOUNCEMENT:**

The company guarantees quality provided that the product is sold in original package.

The damages may be caused by the factors out of the manufacturer's control are at the user's risk.

The company denies all the responsibility for the damages that may result of storing the product under wrong conditions or applying it at wrong doses and times.

## **MANUFACTURER AND LICENSED COMPANY:**

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79