

# 1

AGRI SCIENCES Tarım ve İlaç Ür. San. ve Tic. Ltd. Şti.

# CYMOPRO

## Wetable Powder

Effective Ingredient:70% Propineb + 6% Cymoxanil

Polimeric zinc 1,2-propylrnrbis (dithiocarbamate) + 1-(2-cyano-2-methoxyiminoacetyl)-3-ethylurea

Group:M3-27-Fungicide

**READ THE LABEL FIRST. DO NOT USE AT HOME**

**KEEP AWAY FROM CHILDREN AND FOOD**

**DO NOT INHALE THE VAPOR AND DRIBLETS**

**WEAR MASK, PROTECTIVE UNIFORM, GLOVES, AND GLASSES**

**AVOID SKIN AND EYE CONTACT .**

**DO NOT EAT DURING THE APPLICATION**

**DO NOT SMOKE**

**HUMANS and ANIMALS FOR ARE NOT ALLOWED INTO THE AREA APPLIED FOR 7 DAYS**

**FOLLOW THE INSTRUCTIONS OF USE IN ORDER TO PREVENTS RISKS ON HUMAN and ENVIRONMENT HEALTH**

**STRICTLY PROHIBITED TO USE ON CROPS OTHER THAN THE RECOMMENDED**

**TOXICATION SYMPTOMS:Low toxication probability.**

**FIRST AID PRECAUTIONS:**In case of toxication call the doctor, let him see the product label. Wash well with water and soap in case of contact with body.

ANTIDOTE:No any special antidote. Syptomatic treatment shall be followed.

NATIONAL TOXIC CONSULTANCY ASSOCIATION (NPCA) Tel:114

# 2

Production date:

Expiry date:

Charge nr:

License date and nr:

PLANTS and HAZARDOUS ORGANISMS TO APPLY ON

| Plant     | Savage Herb  | Dose                            | Period between last spraying and harvest |
|-----------|--|---------------------------------|--|
| Zucchini  | False dawning mildew<br>(Pseudoperonospora cubensis) | 50 g/100 L water or<br>200g/da  | 14 days                                  |
| Grapevine | Dawning mildew (plasmopara viticola)                 | 200g/100 L water or<br>200 g/da | 28 days                                  |

## USING DIRECTIONS

### Preparation to Apply:

The advised amount is mixed with some amount of water in a separate tank. The spraying tank is filled in with water in half. While the mixer is on, the mix is added into the tank. The rest of the water is then added into the tank while keeping mixing. This mixing is continued until the spraying is completed.

### Dawny mildew in zucchini :

Spraying must be protective or the latest, when first disease symptoms are seen, and the spraying period should not exceed 7 days. The spraying must be done in the way that each single part of the plants are coated. The sprayings must be continued according to the climate conditions, and the other officially advised products must be used alternately.

### Dawny mildew in grapevine:

First spraying must be done when the sprouts reach to 25-30 cm of length. The second and the third sprayings must be done with periods of 15 days. If the climate conditions are convenient to the movement of the disease.

## 3

Net Amount:

Max. Sales price :

### FACTS OF DANGER:

**Xn:HAZARDOUS**

**N:DANGEROUS FOR NATURE**

### DANGER SYMBOL

.....  
Xn:HAZARDOUS

.....  
N:DANGEROUS FOR NATURE

### INGREDIENTS:

Contains Propineb and Cymoxanil.

### JARGON TO RISK

R20:Hazardous if swallowed

R43:May cause irritation if contacted with skin.

R48/22:Hazardous:Serious damage on health if swallowed continuously for long time.

R51:Very toxic for water organisms

R53:May cause negative effects of long term at water .

## JARGON TO SECURITY

S 2:Keep locked away from children

S13:Keep away from food, beverage, and animal food.

S36/37:Use appropriate protective uniform and gloves

S22:Do not inhale the powder of it

S24:Avoid contact with skin

S46: See the doctor immediately if swallowed, show the package or label

S60: Have this product and its package asided as a hazardous waste.

S46:See the doctor immediately if swallowed, show the package or label

S61:Avoid releasing it into environment. Address to the special use instruction or the Security Information Form.

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79

## 4

### INFORMATION ON RESISTANCE

Cymopro is fungicide classified as Group M3-27 according to its effect mechanism. Repetitive use of products having the same effect mechanism encourages a development of resistance. For this reason, to delay the resistance development, do not exceed the advised total number of applications of Cymopro within a single production period. In cases that requires a repeat of application, it is better to use the plant protect products with different effect mechanism (except Group M3-27)

### MISCIBILITY FACTS

Can be mixed with products that have combination of triadimephone.

## 5

AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.

## CYMOPRO

### Wetable Powder

Effective Ingredient:70% Propineb + 6% Cymoxanil

Polimeric zinc 1,2-propylrnrbis (dithiocarbamate) + 1-(2-cyano-2-methoxyiminoacetyl)-3-ethylurea

**Group:M3-27-Fungicide**

## 6

### USAGE and STORAGE FACTS:

- Keep in its package closed.
- Exterminate the empty packages accordingly.
- Do not use the empty bottle for any other purpose.
- Do not refill the emptied food and beverage packages with the product.
- Toxic for the fish.
- Avoid contact with the underground or surface waters.

### STORING:

Must be stored in its original package, and closed, in a cool place, far away from the food and out of the reach of children, locked. If stored under in a cool and dry place, and in its original pack closed, keeps its physical and chemical specifications min. for two years after its manufacturing date .

### COMPANY'S ANNOUNCEMENT:

Agri Sciences guarantees the quality of this product provided that it is sold in its original package. All the damage that may result of the factors out of the company's control is at the buyer's risk. The company denies all the responsibility for the damages that may result of storing the product under wrong conditions or applying it in wrong amounts or at wrong times.

### **AGRI SCIENCES Tarım ve İlaç Ür. San. Ve Tic. Ltd. Şti.**

Bekir Saydam Cad. No:31,

Pancar Beldesi, Torbalı-İZMİR

Tel:0232 864 13 33, Fax:0232 864 16 79